#### A CATALOGUE OF THE BIRDS OF NORTH AMERICA.

## By ROBERT RIDGWAY.

#### INTRODUCTION.

During the interval of twenty-one years which has elapsed since the publication of the last Smithsonian catalogue,\* a great advance has naturally been made in our knowledge of North American ornithology; and so numerous and important are the changes which have resulted, through additions of new species, rectifications of synonymy, etc., that a new list seems desirable to take the place of the old one.

The total apparent number of species given in the old catalogue has been increased only from 738 to 764, a slight numerical discrepancy which it is necessary to explain. From the catalogue of 1859 there have been eliminated no less than 62 names, which are either not entitled to a place in the North American fauna or which have been degraded to varietal or sub-specific rank, the number of the species in the latter case being here simply duplicated as many times as there are varieties of a species. To offset this large reduction, 59 valid new species have been described since 1859, and 77 added, or restored, to the fauna, the accessions thus numbering 127 species, or 65 more than the eliminations. The forms considered to be of merely subspecific rank number 160, which, added to the 764 valid species recognized, gives a total of 924 definable forms composing the North American avian fauna, as now understood.

It is found impracticable to here distinguish, in all cases, between

TA full analysis of the changes made in this catalogue is given on pages 213-234.

<sup>\*</sup>Two catalogues of North American birds have been issued by the Smithsonian Institution, as follows:

<sup>(1)</sup> Catalogue of North American Birds, chiefly in the Museum of the Smithsonian Institution. By Spencer F. Baird, Assistant Secretary of the Smithsonian Institution. Washington: Smithsonian Institution. October, 1858. 4to, paper. 1 p. l., pp. xviilvi. [Reissue, with new title-page, of pp. xvii-lvi of Vol. IX, Pacific R. R. Reports ("Birds of North America"). Includes, besides the list of 738 species, with habitats, tables of the higher groups, and lists of extralimital species (23 in number) treated in the general report, and of others (31) claimed, on apparently insufficient grounds, as North American; also a summary of the number of species given in the works of Wilson, Bonaparte, and Audubon.

<sup>(2)</sup> Catalogue of North American Birds, chiefly in the Museum of the Smithsonian Institution. By Spencer F. Baird. First octavo edition. Washington: Smithsonian Institution. [Smithsonian Miscellaneous Collections, No. 108.] 1859. Svo. 2 p. ll., pp. 19+2. [Essentially the same as the quarto list, but without habitats, and the matter relating to classification, etc. The two additional pages are an alphabetical index of the North American genera. As in the quarto list, there are, ostensibly, 738 species, but 22 numbers are duplicated, making a total of 760 names in the list.]

species which are truly or peculiarly North American and those which are more properly visitants from other countries: but in the case of those whose occurrence appears to be accidental or occasional, the number preceding the name is inclosed in brackets. Of the latter class, species which there is good reason to believe did not reach our limits through natural means (i. e., those escaped from confinement) have been, in every case, carefully excluded, as have likewise all introduced species.

It has been deemed best, in view of the recent discoveries along our southwestern border, to retain as North American all the species (less than a dozen in number) treated by Professor Baird in Volume IX, Pacific Railroad Reports ("Birds of North America"), and likewise given in the old catalogue, on account of their having been obtained just across the boundary, in Northern Mexico; their discovery within our limits being quite certainly only a question of time and investigation. For the same reason, the remaining few of Giraud's "Sixteen New Species of Texan Birds" are also included. Neither are we prepared to relinquish several Andubonian species which at the present time are known only from the descriptions and figures by their discoverer (e. g., Regulus cuvieri, Perissoglossa [?] carbonata, Dendræca montana, and Wilsonia minuta, as well as other better-known species which are given by Audubon on his own authority (e. g., Chrysomitris "magellanica" = C. notata, and Eudocimus ruber).

Several species peculiar to the islands of Socorro and Guadalupe, off the coast of northwestern Mexico and Lower California, respectively, together with the few forms peculiar to the latter peninsula, are regarded as truly North American, their affinities, with perhaps only two exceptions (i. e., Conurus holochlorus and Polyborus latosus), being strictly "Nearetic."

The greatest difficulty encountered in the compilation of this work has been in the way of distinguishing between valid "species" and those forms to be regarded as geographical races of merely subspecific rank. The greatest care has been taken in all doubtful cases of this kind, and previous conclusions (published in "History of North American Birds"† and elsewhere) carefully reconsidered, with the aid of all the material accessible, including many specimens not previously in hand. This reconsideration of the subject has, in not a few cases, resulted in a reversal of former opinion, specimens from important localities not before represented often deciding the point one way or the other. Every form whose characteristics bear unmistakably the impress of climatic or

<sup>\*</sup>A Description of Sixteen New Species of North American Birds, by Jacob P. Giraud, jr. New York. George F. Nesbitt, printer, Tontine Building, corner of Wall and Water streets. 1841. Folio, not paged, 8 plates. [For species given in this work, which have not since been obtained within the limits of the United States, see p. 229.]

tA History of North American Birds, by S. F. Baird, T. M. Brewer, and R. Ridgway. Land Birds. Illustrated by 64 colored plates and 593 woodcuts. 3 vols., royal 4to. Boston. Little, Brown, & Co.

local influences, gradually less marked toward the habitat of another form, with which it thus intergrades; and all forms which certainly intergrade, no matter how widely distinct the opposite extremes may appear (e. g., Colaptes auratus and mexicanus), together with intergrading forms whose peculiarities are not explained by any known "law" of variation, have been reduced to subspecific rank. On the other hand. where the difference between allied forms is slight, but at the same time absolutely constant, and not coincident with a difference of habitat (e. q., certain of the small Thrushes and the various forms of Junco). specific rank is upheld. There are some forms which future investigation, based upon adequate material, may decide to be of different rank from that accorded them here. We cheerfully acknowledge our fallibility, but at the same time would say that we have endeavored to be as conscientious and consistent as possible, giving the rank of each form as it appears in the light of our present knowledge, independent of previous conclusions.

SMITHSONIAN INSTITUTION, January 22, 1880.

# CATALOGUE.

- 1. HYLOCICHLA MUSTELINA (GMEL.) BAIRD.
  Wood Thrush. [148.]
- 2. HYLOCICHLA FUSCESCENS [STEPH.] BAIRD. Wilson's Thrush. [151.]
- HYLOCICHLA ALICIÆ BAIRD.
   Gray-cheeked Thrush. [154.]
- 4. **HYLOCICHLA USTULATA** (NUTT.) BAIRD. Russet-backed Thrush. [152.]
- 4α. HYLOCICHLA USTULATA SWAINSONI (CABAN.) RIDGW.
  Olive-backed Thrush. [153.]
- 5. HYLOCICHLA UNALASHKÆ (GMEL.) RIDGW.

  Dwarf Thrush. [150.]
- 5a. HYLOCICHLA UNALASHKÆ AUDUBONII (BAIRD) RIDGW.
  Rocky Mountain Hermit Thrush. [149a.]
- 5 b. HYLOCICHLA UNALASHKÆ PALLASI (CABAN.) RIDGW. Hermit Thrush. [149.]
- [6.] TURDUS ILIACUS LINN.

  Red-wing Thrush.
- MERULA MIGRATORIA (LINN.) Sw. & RICH.
   American Robin. [155.]
- 7 a. MERULA MIGRATORIA PROPINQUA RIDGW.
  Western Robin.
- 8. MERULA CONFINIS (BAIRD) RIDGW.
  Saint Lucas Robin.
- 9. HESPEROCICHLA NÆVIA (GMEL.) BAIRD.

  Varied Robin. [156.]
- 10. OREOSCOPTES MONTANUS (TOWNS.) BAIRD. Sage Thrasher. [255.]
- 11. MIMUS POLYGLOTTUS (LINN.) BOIE.

  Mockingbird. [253, 253 a.]

- 12. GALEOSCOPTES CAROLINENSIS (LINN.) CABAN. Catbird. [254.]
- 13. HARPORHYNCHUS RUFUS (LINN.) CABAN. Brown Thrasher. [261, 261 a.]
- 13 a. HARPORHYNCHUS RUFUS LONGIROSTRIS (LAFR.) COUES. Mexican Brown Thrasher, [260.]
- 14. HARPORHYNCHUS CINEREUS XANTUS. Saint Lucas Thrasher.
- 14a. HARPORHYNCHUS CINEREUS BENDIREI (COUES) HENSH. Bendire's Thrasher.
- 15. HARPORHYNCHUS CURVIROSTRIS (Sw.) CABAN. Curve-billed Thrasher. [259, 259 a.]
- 15a, HARPORHYNCHUS CURVIROSTRIS PALMERI RIDGW. Palmer's Thrasher.
- HARPORHYNCHUS REDIVIVUS (GAMB.) CABAN. 16. Californian Thrasher. [256.]
- 16a. HARPORHYNCHUS REDIVIVUS LECONTEI (LAWR.) COUES. Leconte's Thrasher. [257.]
- HARPORHYNCHUS CRISSALIS HENRY. 17. Red-vented Thrasher. [258.]
- 18. HARPORHYNCHUS GRAYSONI BAIRD. Socorro Thrasher.
- 19. CINCLUS MEXICANUS SWAINS. American Water Ouzel. [164.]
- [20.] CYANECULA SUECICA (LINN.) BREHM. Blue-throated Warbler,
- [21.] SAXICOLA CENANTHE (LINN.) BECHST. Stone Chat. [157.]
- 22. SIALIA SIALIS (LINN.) HALDEM Bluebird. [158.]
- 23. SIALIA MEXICANA SWAINS. Californian Bluebird. [159.]
- SIALIA ARCTICA, SWAINS. 24. Rocky Mountain Bluebird. [160.]
- MYIADESTES TOWNSENDI (AUD.) CABAN. 25. Townsend's Solitaire. [235.]
- PHAINOPEPLA NITENS (Sw.) Scl. 26. Black-crested Flycatcher. [234.]
- 27. POLIOPTILA CÆRULEA (LINN.) SCL. Blue-gray Gnatcatcher. [282.]

168	PROCEEDINGS	OF	UNITED	STATES	NATIONAL	MUSEUM.
-----	-------------	----	--------	--------	----------	---------

- 28. POLIOPTILA PLUMBEA BAIRD.

  Plumbeous Gnatcatcher. [283.]
- 29. POLIOPTILA MELANURA LAWR.

  Black-capped Gnatcatcher. [284.]
- 30. REGULUS CALENDULA (LINN.) LICHT.
  Ruby-crowned Kinglet. [161.]
- 31. REGULUS OBSCURUS RIDGW.

  Dusky Kinglet.
- 32. REGULUS CUVIERI Aud.

  Cuvier's Kinglet. [163.]
- 33. · REGULUS SATRAPA LICHT.

  Golden-crowned Kinglet. [162.]
- 33a. REGULUS SATRAPA OLIVACEUS BAIRD.
  Western Golden-crowned Kinglet.
- [34.] PHYLLOSCOPUS BOREALIS (Blas.) Dresser. Kennicott's Warbler.
  - 35. CHAMÆA FASCIATA GAMB.
    Ground Tit. [274.]
  - 36. LOPHOPHANES BICOLOR (Linn.) Bp.
    Tufted Titmouse. [285.]
  - 37. LOPHOPHANES ATROCRISTATUS Cass.

    Black-crested Titmouse. [286.]
  - 38. LOPHOPHANES INORNATUS (GAMB.) CASS.

    Plain Titmouse. [287.]
  - 39. LOPHOPHANES WOLLWEBERI BONAP.
    Wollweber's Titmouse. [288.]
  - 40. PARUS MONTANUS, GAMB.

    Mountain Chickadee. [294.]
  - 41. PARUS ATRICAPILLUS LINN.

    Black-capped Chickadee. [290.]
  - 41a. PARUS ATRICAPILLUS SEPTENTRIONALIS (HARRIS) ALLEN.
    Long-tailed Chickadee. [289, 289a.]
  - 41 b. PARUS ATRICAPILLUS OCCIDENTALIS (BAIRD) COUES.
    Oregon Chickadee. [291.]
  - 42. PARUS CAROLINENSIS Aud.

    Carolina Chickadee. [293.]
  - 43. PARUS MERIDIONALIS Sct.

    Mexican Chickadee. [292.]
  - [44.] PARUS CINCTUS BODD.

    Siberian Chickadee.

- 45. PARUS HUDSONICUS FORST.

  Hudsonian Chickadee, [296.]
- 46. PARUS RUFESCENS Towns.
  Chestnut-backed Chickadee. [295.]
- 46a. PARUS RUFESCENS NEGLECTUS, RIDGW.
  Californian Chickadee.
- 47. PSALTRIPARUS MINIMUS (Towns.) Bp.
  Least Tit. [298.]
- 48. PSALTRIPARUS PLUMBEUS BAIRD.
  Lead-colored Tit. [299.]
- 49. PSALTRIPARUS MELANOTIS (HARTL.) BP.
  Black-eared Tit. [297.]
- 50. AURIPARUS FLAVICEPS (SUNDEV.) BAIRD.
  Yellow-headed Tit. [300.]
- 51. SITTA CAROLINENSIS GMEL.
  White-bellied Nuthatch. [277.]
- 51a. SITTA CAROLINENSIS ACULEATA (CASS.) ALLEN.
  Slender-billed Nuthatch. [278.]
- 52. SITTA CANADENSIS LINN. Red-bellied Nuthatch. [279.]
- SITTA PUSILLA LATH.
   Brown-headed Nuthatch. [280.]
- 54. SITTA PYGMÆA Vig.
  Pigmy Nuthatch. [281.]
- 55. CERTHIA FAMILIARIS RUFA (BARTR.) RIDGW.
  Brown Creeper. [275.]
- 55a. CERTHIA FAMILIARIS MEXICANA (GLOGER) RIDGW.
  Mexican Creeper. [276.]
- 56. CAMPYLORHYNCHUS BRUNNEICAPILLUS (LAFR.) GRAY.
  Cactus Wren. [262.]
- 57. CAMPYLORHYNCHUS AFFINIS XANTUS.
  Saint Lucas Cactus Wren.
- 58. SALPINCTES OBSOLETUS (SAY) CABAN.
  Rock Wren. [264.]
- 58a. SALPINCTES OBSOLETUS GUADALUPENSIS RIDGW.
  Guadalupe Rock Wren.
- 59. CATHERPES MEXICANUS (Sw.) BAIRD.

  Mexican White-throated Wren. [263.]
- 59 a. CATHERPES MEXICANUS CONSPERSUS RIDGW.
  White-throated Wren.

- 170 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
  - 60. THRYOTHORUS LUDOVICIANUS (GM.) Bp. Carolina Wren. [265.].
  - 60 a. THRYOTHORUS LUDOVICIANUS BERLANDIERI (COUCH) COUES.
    Berlandier's Wren. [266.]
  - 60 b. THRYOTHORUS LUDOVICIANUS MIAMENSIS RIDGW. Florida Wren.
  - 61. THRYOMANES BEWICKI (AUD.) BAIRD.

    Bewick's Wren. [267.]
  - 61a. THRYOMANES BEWICKI SPILURUS (Vig.) BAIRD. Californian Bewick's Wren.
  - 61 b. THRYOMANES BEWICKI LEUCOGASTER BAIRD.

    Texan Bewick's Wren.
  - 69. THRYOMANES BREVICAUDA RIDGW.
    Guadalupe Wren.
  - 63. TROGLODYTES AËDON, VIEILL.

    House Wren. [270, 272.]
  - 63a. TROGLODYTES AËDON PARKMANNI (AUD.) COUES. Western House Wren. [271.]
  - 64. TROGLODYTES INSULARIS BAIRD. Socorro Wren.
  - 65. ANORTHURA TROGLODYTES HYEMALIS (VIEILL.) COUES, Winter Wren. [273.]
  - 65 a. ANORTHURA TROGLODYTES PACIFICUS (BAIRD) RIDGW. Western Winter Wren.
  - 66. ANORTHURA ALASCENSIS (BAIRD) COUES.

    Alaskan Wren.
  - 67. TELMATODYTES PALUSTRIS (WILS.) BAIRD.

    Long-billed Marsh Wren. [268.]
  - 67 a. TELMATODYTES PALUSTRIS PALUDICOLA BAIRD Tule Wren.
  - 68. CISTOTHORUS STELLARIS (LICHT.) CABAN. Short-billed Marsh Wren. [269.]
- [69.] MOTACILLA ALBA LINN.
  White Wagtail.
- [70.] BUDYTES FLAVA (LINN.) GRAY.
  Yellow Wagtail.
- 71. ANTHUS LUDOVICIANUS (GM.) LICHT.

  American Titlark. [165.]
- [72.] ANTHUS PRATENSIS (LINN.) BECHST.

  European Titlark.

- 73. NEOCORYS SPRAGUEI (AUD.) Scl. Sprague's Titlark. [166.]
- 74. MNIOTILTA VARIA (LINN.) VIEILL.

  Black-and-white Creeper. [167 a.]
- 74a. MNIOTILTA VARIA BOREALIS (NUTT.) RIDGW.
  Small-billed Creeper. [167.]
- 75. · PROTONOTARIA CITREA (BODD.) BAIRD.
  Prothonotary Warbler. [169.]
- 76. HELONÆA SWAINSONI AUD.

  Swainson's Warbler. [179.]
- 77. HELMITHERUS VERMIVORUS (GMEL.) BP.
  Worm-eating Warbler. [178.]
- 78. HELMINTHOPHAGA BACHMANI (AUD.) CADAN.
  Bachman's Warbler. [182.]
- HELMINTHOPHAGA PINUS (LINN.) BAIRD.
   Blue-winged Yellow Warbler. [180.]
- 80. HELMINTHOPHAGA LAWRENCEI HERRICK.

  Lawrence's Warbler.
- 81. HELMINTHOPHAGA CHRYSOPTERA (LINN.) BAIRD.

  Golden-winged Warbler. [181.]
- 82. HELMINTHOPHAGA LEUCOBRONCHIALIS BREWSTER.
  White-throated Warbler.
- 83. HELMINTHOPHAGA LUCIÆ COOPER.
  Lucy's Warbler.
- 84. HELMINTHOPHAGA VIRGINIÆ BAIRD.

  Virginia's Warbler. [183 a.]
- 85. **HELMINTHOPHAGA RUFICAPILLA** (WILS.) BAIRD.

  Nashville Warbler. [183.]
- 86. HELMINTHOPHAGA CELATA (SAY) BAIRD.
  Orange-crowned Warbler. [184.]
- 86a. HELMINTHOPHAGA CELATA LUTESCENS RIDGW.
  Luteous Warbler.
- 87. HELMINTHOPHAGA PEREGRINA (WILS.) BAIRD
  Tennessee Warbler. [185.]
- 88. PARULA AMERICANA (LINN.) Bp.
  Blue Yellow-backed Warbler. [168.]
- 80. PARULA PITIAYUMI INSULARIS (LAWR.) Ridgw.
  Socorro Warbler.
- 89a. PARULA PITIAYUMI NIGRILORA COUES.
  Sennett's Warbler.

### 172 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.

- 90. PERISSOCIOSSA TIGRINA (GMEL.) BAIRD.

  Cape May Warbler. [206.]
- 91. PERISSOGLOSSA CARBONATA (AUD.) BAIRD. Carbonated Warbler. [207.]
- 92. PEUCEDRAMUS OLIVACEUS (GIRAUD) COUES.
  Olive-headed Warbler.
- 93. DENDRŒCA ÆSTIVA (GMEL.) BAIRD. Summer Yellow Bird. [203.]
- 94. DENDRŒCA CÆRULESCENS (LINN.) BAIRD.

  Black-throated Blue Warbler. [193.]
- 95. DENDRŒCA CORONATA (LINN.) GRAY.
  Yellow-rump Warbler. [194.]
- 96. DENDRŒCA AUDUBONII (Towns.) BAIRD.
  Audubon's Warbler. [195.]
- 97. DENDRŒCA MACULOSA (GMEL.) BAIRD.

  Black-and-yellow Warbler. [204.]
- 98. DENDRŒCA CÆRULEA (Wils.) BAIRD. Cerulean Warbler. [201.]
- 99. **DENDRŒCA PENNSYLVANICA** (Linn.) Baird. Chestnut-sided Warbler. [200.]
- 100. DENDRŒCA CASTANEA (WILS.) BAIRD. Bay-breasted Warbler. [197.]
- 101. DENDRŒCA STRIATA (FORST.) BAIRD.
  Black-poll Warbler. [202.]
- 102. DENDRŒCA BLACKBURNIÆ (Gm.) BAIRD.
  Blackburnian Warbler. [196.]
- 103. DENDRŒCA DOMINICA (LINN.) BAIRD.
  Yellow-throated Warbler. [209.]
- 103 a. DENDRŒCA DOMINICA ALBILORA BAIRD.

  White-browed Yellow-throated Warbler,
- 104. DENDRŒCA GRACIÆ COUES.

  Grace's Warbler.
- 105. DENDRŒCA NIGRESCENS (TOWNS.) BAIRD. Black-throated Gray Warbler. [192.]
- 106. DENDRŒCA CHRYSOPARIA Scl. & Salv. Golden-cheeked Warbler.
- 107. DENDRŒCA VIRENS (GMEL.) BAIRD.

  Black-throated Green Warbler. [189.]
- 103. DENDRŒCA TOWNSENDI (NUTT.) BAIRD. Townsend's Warbler. [191.]

- 109. DENDRŒCA OCCIDENTALIS (Towns.) BAIRD.

  Hermit Warbler. [190.]
- 110. DENDRŒCA KIRTLANDI BAIRD. Kirtland's Warbler. [205.]
- 111. DENDRŒCA PINUS (WILS.) BAIRD.
  Pine-creeping Warbler. [198.]
- 112. DENDRŒCA MONTANA (WILS.) BAIRD.

  Blue Mountain Warbler. [199.]
- 113. **DENDRŒCA PALMARUM** (GMEL.) BAIRD. Red-poll Warbler. [208.]
- 113a, DENDRŒCA PALMARUM HYPOCHRYSEA RIDGW.
  Yellow Red-poll Warbler.
- 114. DENDRŒCA DISCOLOR (VIEILL.) BAIRD.

  Prairie Warbler. [210.]
- 115. SIURUS AURICAPILLUS (LINN.) SWAINS.

  Golden-crowned Thrush. [186.]
- 116. SIURUS NÆVIUS (BODD.) COUES.
  Small-billed Water Thrush. [187.]
- 116a. SIURUS NÆVIUS NOTABILIS GRINNELL.
  Grinnell's Water Thrush.
- 117. SIURUS MOTACILLA (VIEILL.) Coues.

  Large-billed Water Thrush. [188.]
- 118. OPORORNIS AGILIS (WILS.) BAIRD.

  Connecticut Warbler. [174.]
- 119. OPORORNIS FORMOSA (Wils.) BAIRD. Kentucky Warbler. [175.]
- 120. GEOTHLYPIS PHILADELPHIA (Wils.) BAIRD.

  Mourning Warbler. [172.]
- 121. **GEOTHLYPIS MACGILLIVRAYI** (AUD.) BAIRD. **Macgillivray**'s Warbler. [173.]
- 122. GEOTHLYPIS TRICHAS (LINN.) CABAN.

  Maryland Yellow-throat. [170.]
- 123. ICTERIA VIRENS (LINN.) BAIRD.

  Yellow-breasted Chat. [176.]
- 123a. ICTERIA VIRENS LONGICAUDA (LAWR.) COUES.
  Long-tailed Chat. [177.]
- 124. WILSONIA MITRATA (GMEL.) Bp.

  Hooded Warbler, [211.]
- 125. WILSONIA PUSILLA (WILS.) BP. Black-capped Yellow Warbler. [213.]

- 174 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 125 a. WILSONIA PUSILLA PILEOLATA (PALL.) RIDGW. Pileolated Warbler. .
- 126. WILSONIA MINUTA (WILS.) RIDGW.
  Small-headed Flycatcher. [212.]
- 127. WILSONIA CANADENSIS (LINN.) COUES.

  Canadian Flycatching Warbler. [214, 215.]
- 128. SETOPHAGA RUTICILLA (LINN.) SWAINS.

  American Redstart. [217.]
- 129. SETOPHAGA PICTA SWAINS.

  Painted Redstart. [218.]
- 130. SETOPHAGA MINIATA SWAINS.
  Red-bellied Redstart. [919]
- 131. CARDELLINA RUBRIFRONS (GIRAUD) Scl.
  Red-faced Warbler.
- 132. ERGATICUS RUBER (SWAINS.) BAIRD.
  Red Warbler. [216.]
- 133. BASILEUTERUS CULICIVORUS (LICHT.) BONAP.

  Brasier's Warbler.
- 134. BASILEUTERUS BELLII (GIRAUD) SCL. Bell's Warbler.
- 135. VIREOSYLVIA OLIVACEA (LINN.) BONAP. Red-eyed Vireo. [240.]
- 136. VIREOSYLVIA AGILIS FLAVO-VIRIDIS (CASS.) RIDGW.
  Yellow-green Vireo. [241.]
- 137. VIREOSYLVIA CARIDRIS BARBATULA (CABAN.) RIDGW.
  Black-whiskered Vireo. [243.]
- 138. VIREOSYLVIA PHILADELPHICA CASS.
  Philadelphia Vireo. [244.]
- 139. VIREOSYLVIA GILVA (VIEILL.) CASS.
  Warbling Vireo. [245.]
- 139 a. VIREOSYLVIA GILVA SWAINSONI BAIRD.
  Western Warbling Vireo.
- 140. LANIVIREO FLAVIFRONS (VIEILL.) BAIRD.
  Yellow-throated Vireo. [252.]
- 141. LANIVIREO SOLITARIUS (VIEILL.) BAIRD.

  Blue-headed Vireo. [250.]
- 141 a. LANIVIREO SOLITARIUS CASSINI (XANTUS) RIDGW.

  Cassin's Vireo. [251.]
- 141 b. LANIVIREO SOLITARIUS PLUMBEUS (COUES) ALL. Plumbeous Vireo.

- 142. VIREO ATRICAPILLUS WOODH.

  Black-capped Vireo. [247.]
- 143. VIREO NOVEBORACENSIS (GMEL.) Bp.
  White-eyed Vireo. [248.]
- 144. VIREO HUTTONI CASS.

  Hutton's Vireo. [249.]
- 145. VIREO BELLII AUD.

  Bell's Vireo. [246.]
- 146. VIREO PUSILLUS COUES.

  Least Vireo.
- 147. VIREO VICINIOR COUES.

  Gray Vireo.
- 143. LANIUS BOREALIS VIEILL.

  Great Northern Shrike. [236.]
- 149. LANIUS LUDOVICIANUS LINN.
  Loggerhead Shrike. [237.]
- 149a. LANIUS LUDOVICIANUS EXCUBITORIDES (Sw.) Coues.
  White-rumped Shrike. [233.]
- 149 b. LANIUS LUDOVICIANUS ROBUSTUS BAIRD.

  Large-billed Shrike.
- 150. AMPELIS GARRULUS LINN.
  Northern Wax-wing. [232.]
- 151. AMPELIS CEDRORUM (VIEILL.) BAIRD.
  Cedar Wax-wing. [233.]
- 152. PROGNE SUBIS (LINN.) BAIRD.

  Purple Martin. [231.]
- 152 a. PROGNE SUBIS CRYPTOLEUCA BAIRD.

  Cuban Martin. [231 a.]
- 153. **PETROCHELIDON LUNIFRONS** (SAY) LAWR. Cliff Swallow. [226.]
- 154. HIRUNDO ERYTHROGASTRA BODD.

  Barn Swallow. [925.]
- 155. TACHYCINETA BICOLOR (VIEILL.) CABAN.
  White-bellied Swallow. [227.]
- 156. TACHYCINETA THALASSINA (SWAINS.) CABAN.
  Violet-green Swallow. [228.]
- 157. COTILE RIPARIA (LINN.) BOIE.

  Bank Swallow. [229.]
- 158. STELGIDOPTERYX SERRIPENNIS (AUD.) BAIRD.
  Rough-winged Swallow. [230.]

- 176 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 159. CERTHIOLA BAHAMENSIS REICHENB.

  Bahaman Honey Creeper. [301.]
- 160. EUPHONIA ELEGANTISSIMA (Bp.) Gray.
  Blue-headed Euphonia. [224.]
- 161. PYRANGA RUBRA (LINN.) VIEILL.
  Scarlet Tanager. [220.]
- 162. PYRANGA LUDOVICIANA (Wils.) Bp. Western Tanager. [223.]
- 163. PYRANGA HEPATICA SWAINS.

  Hepatic Tanager. [222.]
- 164. PYRANGA ÆSTIVA (LINN.) VIEILL. Summer Redbird. [221.]
- 164 a. PYRANGA ÆSTIVA COOPERI RIDGW. Cooper's Tanager.
- 165. **HESPERIPHONA VESPERTINA** (COOPER) Bp. Evening Grosbeak. [303.]
- 166. PINICOLA ENUCLEATOR (LINN.) VIEILL.
  Pine Grosbeak. [304.]
- [167.] PYRRHULA CASSINI BAIRD.

  Alaskan Bullfinch.
- 168. CARPODACUS PURPUREUS (GM.) BAIRD.
  Purple Finch. [305.]
- 168a. CARPODACUS PURPUREUS CALIFORNICUS BAIRD. Californian Purple Finch. [306.]
- 169. CARPODACUS CASSINI BAIRD.

  Cassin's Purple Finch. [307.]
- 170. CARPODACUS FRONTALIS (SAY) GRAY.
  House Finch. [308.]
- 170 a. CARPODACUS FRONTALIS RHODOCOLPUS (CABAN.) RIDGW.
  Crimson House Finch.
- 171. CARPODACUS AMPLUS RIDGW.
  Guadalupe House Finch.
- 172. LOXIA CURVIROSTRA AMERICANA (WILS.) COUES.
  American Crossbill. [318.]
- 172 a. LOXIA CURVIROSTRA MEXICANA (STRICKL.) BAIRD.

  Mexican Crossbill. [318 a.]
- 173. LOXIA LEUCOPTERA GM.
  White-winged Crossbill. [319.]
- 174. LEUCOSTICTE GRISEINUCHA (BRANDT) BAIRD.
  Aleutian Rosy Finch. [323.]

- 175. LEUCOSTICTE TEPHROCOTIS SWAINS.

  Gray-crowned Rosy Finch. [322.]
- 175 a. LEUCOSTICTE TEPHROCOTIS LITTORALIS (BAIRD) COUES.
  Gray-headed Rosy Finch.
- 176. LEUCOSTICTE ATRATA RIDGW.

  Black Rosy Finch.
- 177. LEUCOSTICTE AUSTRALIS ALLEN.
  Brown-capped Rosy Finch.
- 178. ÆGIOTHUS CANESCENS GOULD.
  Mealy Redpoll. [321.]
- 178a. ÆGIOTHUS CANESCENS EXILIPES (COUES) RIDGW.
  White-rumped Redpoll.
- 179. ÆGIOTHUS LINARIA (LINN.) CABAN.

  Common Redpoll. [320.]
- 179a. ÆGIOTHUS LINARIA HOLBOLLI (BREHM) RIDGW.
  Greater Redpoll.
- 180. ÆGIOTHUS BREWSTERI RIDGW.
  Brewster's Linnet.
- 181. ASTRAGALINUS TRISTIS (LINN.) CAB.
  American Goldfinch. [313.]
- 182. ASTRAGALINUS PSALTRIA (SAY) COUES.
  Green-backed Goldfinch. [314.]
- 182a. ASTRAGALINUS PSALTRIA ARIZONÆ, COUES.
  Arizona Goldfinch. [315.]
- 182 b. ASTRAGALINUS PSALTRIA MEXICANUS (Sw.) Coues.

  Mexican Goldfinch.
- 183. ASTRAGALINUS LAWRENCEII (CASS.) Bp. Lawrence's Goldfinch. [316.]
- 184. CHRYSOMITRIS NOTATA (Du Bus) Bp.
  Black-headed Goldfinch. [310.]
- 185. CHRYSOMITRIS PINUS (WILS.) BP.
  Pine Goldfinch. [317.]
- 186. PLECTROPHANES NIVALIS (LINN.) MEYER.
  Snow Bunting. [325.]
- 187. CENTROPHANES LAPPONICUS (LINN.) CABAN.
  Lapland Longspur. [326.]
- 188. \*CENTROPHANES PICTUS (SWAINS.) CABAN.
  Smith's Longspur. [327.]
- 189. CENTROPHANES ORNATUS (TOWNS.) CABAN.

  Chestnut-collared Longspur. [328, 329.]

  Proc. Nat. Mus. 80——12

  Aug. 24, 1880.

17	78	PROC	FEDINGS	OF III	CITED	STATES	NATIONAL	MIISPIIM
	0	LUOU	PEDINGS	OF U	NITED	BITTER	MATIONAL	DI COL CUI.

- 190. RHYNCHOPHANES MACCOWNI (LAWR.) BAIRD McCown's Longspur. [330.]
- 191. CENTRONYX BAIRDII (Aud.) BAIRD.

  Baird's Bunting. [331.]
- 192. PASSERCULUS PRINCEPS MAYNARD.

  Ipswich Sparrow.
- 193. PASSERCULUS SANDWICHENSIS (GMEL.) BAIRD. Sandwich Sound Sparrow. [333.]
- 193 a. PASSERCULUS SANDWICHENSIS SAVANNA (WILS.) RIDGW. Savannah Sparrow. [332.]
- 193b. PASSERCULUS SANDWICHENSIS ALAUDINUS (Bp.) RIDGW Western Savannah Sparrow. [335.]
- 194. PASSERCULUS ANTHINUS BONAP.
  Titlark Sparrow. [334.]
- 195. PASSERCULUS GUTTATUS LAWR.
  Saint Lucas Sparrow.
- 196. PASSERCULUS ROSTRATUS (CASS.) BAIRD.

  Large-billed Sparrow. [336.]
- 197. POŒCETES GRAMINEUS (GM.) BAIRD.
  Grass Finch. [337.]
- 107 a. POŒCETES GRAMINEUS CONFINIS BAIRD. Western Grass Finch.
- 193. COTURNICULUS PASSERINUS (WILS.) Bp. Yellow-winged Sparrow. [338.]
- 198 a. COTURNICULUS PASSERINUS PERPALLIDUS RIDGW.
  Western Yellow-winged Sparrow
- 199. COTURNICULUS HENSLOWI (AUD.) BP. Henslow's Sparrow. [339.]
- 200. COTURNICULUS LECONTEI (AUD.) Bp. Leconte's Sparrow. [340.]
- 201. AMMODROMUS CAUDACUTUS (Gm.) Swains.
  Sharp-tailed Finch. [341.]
- 201a. AMMODROMUS CAUDACUTUS NELSONI ALLEN.
  Nelson's Sharp-tailed Finch.
- 202. AMMODROMUS MARITIMUS (WILS.) SWAINS.
  Sea-side Finch. [342.]
- 203. AMMODROMUS NIGRESCENS RIDGW. Black-and-white Sea-side Finch.
- 204. CHONDESTES GRAMMICA (SAY) Bp. Lark Finch. [344.]

- 204a. CHONDESTES GRAMMICA STRIGATA (Sw.) RIDGW.
  Western Lark Finch.
- 205. ZONOTRICHIA QUERULA (NUTT.) GAMB. Harris's Sparrow. [348.]
- 206. ZONOTRICHIA LEUCOPHRYS (FORST.) SWAINS.
  White-crowned Sparrow. [345.]
- 207. ZONOTRICHIA GAMBELI (NUTT.) GAMB.
  Gambel's White-crowned Sparrow.
- 207 a. ZONOTRICHIA GAMBELI INTERMEDIA RIDGW.

  Intermediate White-crowned Sparrow. [346,]
- 203. ZONOTRICHIA CORONATA (PALL.) BAIRD.
  Golden-crowned Sparrow. [347.]
- 209. ZONOTRICHIA ALBICOLLIS (GM.) Bp. White-throated Sparrow. [349.]
- 210. SPIZELLA MONTANA (FORST.) RIDGW.

  Tree Sparrow. [357.]
- 211. SPIZELLA DOMESTICA (BARTR.) COUES.
  Chipping Sparrow. [359.]
- 211a. SPIZELLA DOMESTICA ARIZONÆ (Coues) Ridgw.
  Western Chipping Sparrow.
- 212. SPIZELLA PALLIDA (Sw.) Bp.
  Clay-colored Sparrow. [360.]
- 213. SPIZELLA BREWERI Cass.

  Brewer's Sparrow. [361.]
- 214. SPIZELLA PUSILLA (WILS.) BP. Field Sparrow. [358.]
- 215. SPIZELLA ATROGULARIS (CABAN.) Bb.
  Black-chinned Sparrow. [362.]
- 216. JUNCO AIKENI RIDGW.

  White-winged Snowbird.
- 217. JUNCO HYEMALIS (LINN.) Scl.
  Black Snowbird. [354.]
- 218. JUNCO OREGONUS (Towns.) Scl., Oregon Snowbird. [352.]
- 219. JUNCO ANNECTENS BAIRD.

  Pink-sided Snowbird.
- 220. JUNCO CANICEPS (WOODH.) BAIRD.

  Gray-headed Snowbird. [353.]
- 221. JUNCO DORSALIS HENRY.

  Red-backed Snowbird. [351.]

- 180 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 222. JUNCO CINEREUS (SWAINS.) CABAN.

  Mexican Snowbird. [350.]
- 223. JUNCO INSULARIS RIDGW.
  Guadalupe Snowbird.
- 224. AMPHISPIZA BILINEATA (Cass.) Coues.

  Black-throated Sparrow. [355.]
- 225. AMPHISPIZA BELLII (Cass.) Coues. Bell's Sparrow. [356.]
- 225 a. AMPHISPIZA BELLII NEVADENSIS RIDGW. Sagebrush Sparrow.
- 226. PEUCÆA ÆSTIVALIS (LICHT.) CABAN. Bachman's Finch. [370.]
- 226 a. PEUCÆA ÆSTIVALIS ILLINOENSIS RIDGW.
  Oak-woods Sparrow.
- 227. PEUCÆA ARIZONÆ RIDGW.
  Arizona Sparrow.
- 228. PEUCÆA CASSINI (WOODH.) BAIRD.

  Cassin's Sparrow. [371.]
- 229. PEUCÆA CARPALIS COUES.

  Rufous-winged Sparrow.
- 230. PEUCÆA RUFICEPS (Cass.) Baird.
  Rufous-crowned Sparrow. [372.]
- 230 a. PEUCÆA RUFICEPS BOUCARDI (Scl.) B. B. & R. Boucard's Sparrow.
- 231. MELOSPIZA FASCIATA (FORST.) SCOTT. Song Sparrow. [363.]
- 231 a. MELOSPIZA FASCIATA FALLAX BAIRD.

  Mountain Song Sparrow. [367.]
- 231 b. MELOSPIZA FASCIATA HEERMANNI BAIRD. Heermann's Song Sparrow. [364.]
- 231 c. MELOSPIZA FASCIATA SAMUELIS BAIRD.

  Californian Song Sparrow. [343, 365.]
- 231 d. MELOSPIZA FASCIATA GUTTATA (NUTT.) BAIRD.
  Rusty Song Sparrow. [366.]
- 231 c. MELOSPIZA FASCIATA RUFINA (BRANDT) BAIRD. Sooty Song Sparrow.
- 232. MELOSPIZA CINEREA (GM.) RIDGW.
  Aleutian Song Sparrow.
- 233. MELOSPIZA PALUSTRIS (WILS.) BAIRD. Swamp Sparrow. [369.]

- 234. MELOSPIZA LINCOLNI (AUD.) BAIRD. Lincoln's Finch. [368.]
- 235. PASSERELLA ILIACA (MERREM) Sw.
  Fox-colored Sparrow. [374.]
- 235 a. PASSERELLA ILIACA UNALASHKENSIS (Gm.) RIDGW.
  Townsend's Sparrow, [375.]
- 235 b. PASSERELLA ILIACA MEGARHYNCHA (BAIRD) RIDGW.
  Thick-billed Sparrow. [376a.]
- 235 c. PASSERELLA ILIACA SCHISTACEA (BAIRD) ALLEN.
  Slate-colored Sparrow. [376.]
- 236. EMBERNAGRA RUFIVIRGATA LAWR.
  Texas Sparrow. [373.]
- 237. PIPILO ERYTHROPHTHALMUS (LINN.) VIEILL.
  Chewink; Towhee. [391.]
- 237 a. PIPILO ERYTHROPHTHALMUS ALLENI COUES.
  Florida Towhee.
- 238. PIPILO MACULATUS ARCTICUS (SWAINS.) COUES.

  Northern Towhee. [393.]
- 238a. PIPILO MACULATUS MEGALONYX (BAIRD) COUES.

  Spurred Towhee. [394.]
- 238b. PIPILO MACULATUS OREGONUS (Bell) Coues. Oregon Towhee. [392.]
- 238 c. PIPILO MACULATUS CONSOBRINUS RIDGW.
  Guadalupe Towhee.
- 238d. PIPILO MACULATUS CARMANI BAIRD.
  Socoito Towhee.
- 239. PIPILO CHLORURUS (Towns.) Baird.

  Green-tailed Towhee. [398.]
- 240. PIPILO FUSCUS MESOLEUCUS (BAIRD) RIDGW.

  Cañon Towhee. [397.]
- 240 a. PIPILO FUSCUS ALBIGULA (BAIRD) COUES.

  Saint Lucas Brown Towhee.
- 240 b. PIPILO FUSCUS CRISSALIS (Vig.) Coues.

  Californian Brown Towhee. [396.]
- 241 PIPILO ABERTI BAIRD.

  Abert's Towhee. [395.]
- 242. CARDINALIS VIRGINIANUS (BRISS.) BP.
  Cardinal Grosbeak. [390.]
- 242 a. CARDINALIS VIRGINIANUS IGNEUS (BAIRD) COUES.
  Saint Lucas Cardinal.

- 182 PROCEEDINGS OF UNITED STATES. NATIONAL MUSEUM.
- 243. PYRRHULOXIA SINUATA BONAP.

  Texan Cardinal. [389.]
- 244. ZAMELODIA LUDOVICIANA (LINN.) COUES.

  Rose-breasted Grosbeak. [380.]
- 245. ZAMELODIA MELANOCEPHALA (SWAINS.) COUES.

  Black-headed Grosbeak. [381.]
- 246. GUIRACA CÆRULEA (LINN.), SWAINS.

  Blue Grosbeak. [382.]
- 247. PASSERINA PARELLINA (Bp.) RIDGW. Blue Bunting. [383.]
- 248. PASSERINA CYANEA (LINN.) GRAY.
  Indigo Bunting. [387.]
- 249. PASSERINA AMŒNA (SAY) GRAY. Lazuli Bunting. [386.]
- 250. PASSERINA VERSICOLOR (BONAP.) GRAY.
  Varied Bunting. [385.]
- 251. PASSERINA CIRIS (LINN.) GRAY.
  Painted Bunting; Nonpareil. [384.]
- 259. SPERMOPHILA MORELETII PUCHERAN.

  Morelet's Seedeater. [388.]
- 253. PHONIPARA ZENA (LINN.) BRYANT.
  Black-faced Seedeater.
- 254. SPIZA AMERICANA (Gm.) BONAP.

  Black-throated Bunting. [378.]
- 255. SPIZA TOWNSENDI (AUD.) RIDGW.
  Townsend's Bunting. [379.]
- 256. CALAMOSPIZA BICOLOR (TOWNS.) BONAP.

  Lark Bunting. [377.]
- 257. DOLICHONYX ORYZIVORUS (LINN.) SWAINS. Bobolink. [399.]
- 258. MOLOTHRUS ATER (BODD.) GRAY.

  Cowbird. [400.]
- 258 a. MOLOTHRUS ATER OBSCURUS (GMEL.) COUES.

  Dwarf Cowbird.
- 259. MOLOTHRUS ÆNEUS (WAGL.) CABAN. Bronzed Cowbird.
- 260. XANTHOCEPHALUS ICTEROCEPHALUS (BONAP.) BD. Yellow-headed Blackbird. [404.]
- 261. AGELÆUS PHŒNICEUS (LINN.) VIEILL.

  Red-and-buff-shouldered Blackbird. [401.]

- 261a. AGELÆUS PHŒNICEUS GUBERNATOR (WAGL.) COUES.
  Red-and-black-shouldered Blackbird. [402.]
- 262. AGELÆUS TRICOLOR (NUTT.) BP.

  Red-and-white-shouldered Blackbird. [403.]
- 263. STURNELLA MAGNA (LINN.) SWAINS.

  Meadow Lark. [406.]
- 263 a. STURNELLA MAGNA MEXICANA (Scl.) RIDGW.
  Mexican Meadow Lark.
- 264. STURNELLA NEGLECTA AUD.

  Western Meadow Lark. [407.]
- 205. ICTERUS VULGARIS DAUD. Troupial. [408.]
- 266. ICTERUS AUDUBONII GIRAUD.

  Audubon's Oriole. [409.]
- 967. ICTERUS WAGLERI Scl.
  Wagler's Oriole. [412.]
- Wagler's Oriole. [412.]
  268. ICTERUS PARISORUM BONAP.
- 269, ICTERUS CUCULLATUS SWAINS.

  Hooded Oriole. [413.]
- 270. ICTERUS SPURIUS (LINN.) Bp.
  Orchard Oriole. [414.]
- 271. ICTERUS GALBULA (LINN.) COUES.

  Baltimore Oriole. [415.]
- 272. ICTERUS BULLOCKI (SWAINS.) Bp. Bullock's Oricle. [416.]
- 273. SCOLECOPHAGUS FERRUGINEUS (GM.) SWAINS.
  Rusty Blackbird. [417.]

Scott's Oriole. [411.]

- 274. SCOLECOPHAGUS CYANOCEPHALUS (WAGL.) CABAN.
  Brewer's Blackbird. [418.]
- 275. QUISCALUS MACRURUS SWAINS.

  Great-tailed Grackle. [419.]
- 276. QUISCALUS PALUSTRIS SWAINS.
  Mexican Boat-tailed Grackle.
- 277. QUISCALUS MAJOR VIEILL.

  Boat-tailed Grackle. [420.]
- 278. QUISCALUS PURPUREUS (BARTR.) LEICHT.
  Purple Grackle. [421.]
- 278α. QUISCALUS PURPUREUS AGLÆUS (BAIRD) COUES. Florida Grackle. [402.]

- 184 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 278 b. QUISCALUS PURPUREUS ÆNEUS RIDGW. Bronzed Grackle.
- [279.] STURNUS VULGARIS LINN.

  European Starling.
- 280. CORVUS CORAX CARNIVORUS (BARTR.) RIDGW.

  American Raven. [423, 424.]
- 281. CORVUS CRYPTOLEUCUS COUCH.
  White-necked Raven. [425.]
- 282. CORVUS FRUGIVORUS BARTR.

  Common Crow. [426.]
- 282 a. CORVUS FRUGIVORUS FLORIDANUS (BAIRD) RIDGW. Florida Crow. [427.]
- 282 b. CORVUS FRUGIVORUS CAURINUS (BAIRD) RIDGW.
  Northwestern Fish Crow. [428.]
- 283. CORVUS OSSIFRAGUS WILS. Fish Crow. [429.]
- 284. PICICORVUS COLUMBIANUS (Wils.) Bp. Clarke's Nutcracker. [430.]
- 285. GYMNOCITTA CYANOCEPHALA Max.

  Maximilian's Nutoracker; Piñon Jay. [431.]
- 286. PICA RUSTICA HUDSONICA (Scop.) BAIRD.
  Black-billed Magpie. [432.]
- 287. PICA NUTTALLI Aud.
  Yellow-billed Magpie. [433.]
- 288. PSILORHINUS MORIO (WAGL.) GRAY. Brown Jay. [444.]
- 289. CYANOCITTA CRISTATA (LINN.) STRICKL.
  Blue Jay. [434.]
- 290. CYANOCITTA STELLERI (GM.) CABAN. Steller's Jay. [435.]
- 290 a. CYANOCITTA STELLERI FRONTALIS RIDGW. Blue-fronted Jay.
- 290 b. CYANOCITTA STELLERI ANNECTENS (BAIRD) RIDGW. Black-headed Jay.
- 290 c. CYANOCITTA STELLERI MACROLOPHA (BAIRD) RIDGW.,
  Long-crested Jay. [436.]
- 291. APHELOCOMA FLORIDANA (BARTR.) CABAN. Florida Jay. [439.]
- 292. APHELOCOMA WOODHOUSEI (BAIRD) RIDGW. Woodhouse's Jay. [438.]

- 293. APHELOCOMA CALIFORNICA (Vig.) CABAN, California Jay. [437.]
- 294. APHELOCOMA ULTRAMARINA COUCHII BAIRD. Couch's Jay. [441.]
- 295. APHELOCOMA SORDIDA ARIZONÆ RIDGW.
  Arizona Jay. [440.]
- 296. XANTHURA LUXUOSA (LESS.) BP. Green Jay. [442.]
- 297. PERISOREUS CANADENSIS (LINN.) Bp.
  Canada Jay. [443.]
- 297 a. PERISOREUS CANADENSIS CAPITALIS BAIRD.
  White-headed Jay.
- 297 b. PERISOREUS CANADENSIS FUMIFRONS RIDGW. Smoky-fronted Jay.
- 298. PERISOREUS OBSCURUS RIDGW.
  Oregon Jay.
- [299.] ALAUDA ARVENSIS LINN. Sky Lark.
- 300. EREMOPHILA ALPESTRIS (FORST.) BOIE.

  Shore Lark. [302.]
- 300 a. EREMOPHILA ALPESTRIS LEUCOLÆMA Coues.
  White-throated Shore Lark.
- 300 b. EREMOPHILA ALPESTRIS CHRYSOLÆMA (WAGL.) COUES.

  Mexican Shore Lark.
- 301. MILVULUS FORFICATUS (GM.) SWAINS:
  Scissor-tailed Flycatcher. [123.]
- [302.] MILVULUS TYRANNUS (LINN.) Bp.

  Fork-tailed Flycatcher. [122.]
- 303. TYRANNUS DOMINICENSIS (GM.) REICH.
  Gray Kingbird. [125.]
- 304. TYRANNUS CAROLINENSIS (LINN.) TEMM.
  Kingbird; Bee Martin. [124.]
- 305. TYRANNUS MELANCHOLICUS COUCHII BAIRD.
  Couch's Kingbird. [128, 129.]
- 306. TYRANNUS VERTICALIS SAY.
  Western Kingbird. [126.]
- 307. TYRANNUS VOCIFERANS SWAINS.

  Cassin's Kingbird. [127.]
- 308. PITANG**US** DERBIANUS (KAUP) Scl. Mexican Pitangus.

- 186 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 309. MYIOZETETES TEXENSIS (GIRAUD) Scl. Giraud's Flycatcher.
- 310. MYIODINASTES LUTEIVENTRIS BONAP.
  Henshaw's Flycatcher.
- 311. MYIARCHUS MEXICANUS (KAUP) LAWR.

  Mexican Great Crested Flycatcher. [132.]
- 312. MYIARCHUS CRINITUS (LINN.) CABAN,
  Great Crested Flycatcher. [130.]
- 313. MYIARCHUS CINERASCENS LAWR.

  Ash-throated Flycatcher. [131.]
- 314. MYIARCHUS LAWRENCEI (GIRAUD) BAIRD.

  Lawrence's Flycatcher. [133.]
- 315. SAYORNIS FUSCUS (GMEL.) BAIRD.

  Phœbe Bird; Pewee. [135.]
- 316. SAYORNIS SAYI (BONAP.) BAIRD. Say's Pewee. [136.]
- 317. SAYORNIS NIGRICANS (SWAINS.) BP. Black Pewee. [134.]
- 318. CONTOPUS BOREALIS (SWAINS.) BAIRD.
  Olive-sided Flycatcher. [137.]
- 319. CONTOPUS PERTINAX CABAN.
  Coues's Flycatcher.
- 320. CONTOPUS VIRENS (LINN.) CABAN.
  Wood Pewee. [139.]
- 321. CONTOPUS RICHARDSONII (Sw.) BAIRD.
  Western Wood Pewee. [133.]
- 322. EMPIDONAX FLAVIVENTRIS BAIRD.
  Yellow-bellied Flycatcher. [144.]
- 323. EMPIDONAX DIFFICILIS BAIRD.

  Western Yellow-bellied Flycatcher. [144a.]
- 324. EMPIDONAX ACADICUS (GMEL.) BAIRD.
  Acadian Flycatcher. [143.]
- 325. EMPIDONAX PUSILLUS (SWAINS.) BD. Little Flycatcher. [141.]
- 325 a. EMPIDONAX PUSILLUS TRAILLII (AUD.) BAIRD.
  Traill's Flycatcher. [140.]
- 326. EMPIDONAX MINIMUS BAIRD.

  Least Flycatcher. [142.]
- 327. EMPIDONAX HAMMONDI (XANTUS) BD. Hammond's Flycatcher. [145.]

- 328. EMPIDONAX OBSCURUS (SWAINS.) BAIRD.
  Wright's Flycatcher. [146.]
- 329. EMPIDONAX FULVIFRONS (GIRAUD) Scl. Fulvous Flycatcher.
- 329 a. EMPIDONAX FULVIFRONS PALLESCENS Coues.

  Buff-breasted Flycatcher.
- 330. PYROCEPHALUS RUBINEUS MEXICANUS (Scl.) Cours.

  Vermilion Flycatcher. [147.]
- 331. ORNITHION IMBERBE (Scl.) Cours.
  Small-billed Flycatcher.
- 332. PACHYRHAMPHUS MAJOR (BONAP.) Scl., Thick-billed Flycatcher. [121.]
- 333. HADROSTOMUS AGLAIÆ (LAFR.) CAB.

  Rose-throated Flycatcher. [120.]
- 334. EUGENES FULGENS (SWAINS.) GOULD.
  Refulgent Hummingbird.
- 335. TROCHILUS COLUBRIS LINN.

  Ruby-throated Hummingbird. [101.]
- 336. TROCHILUS ALEXANDRI BOURC. & MULS.

  Black-chinned Hummingbird. [102.]
- 337. CALYPTE COSTÆ (BOURC.) GOULD.

  Costa's Hummingbird. [106]
- 338. CALYPTE ANNÆ (LESS.) GOULD.

  Anna's Hummingbird. [105.]
- 339. SELASPHORUS PLATYCERCUS (Swåins.) Bp. Broad-tailed Hummingbird. [104.]
- 340. **SELASPHORUS RUFUS** (GMEL.) AUD. Rufous Hummingbird. [103.]
- 341. SELASPHORUS ALLENI HENSH.
  Allen's Hummingbird.
- 342. ATTHIS HELOISÆ (LESS.) REICH.
  Heloise's Hummingbird.
- 343. STELLULA CALLIOPE GOULD.

  Calliope Hummingbird.
- 344. CALOTHORAX LUCIFER (SWAINS.) GRAY. Lucifer Hummingbird.
- 345. AMAZILIA FUSCICAUDATA (FRASER) RIDGW. Rieffer's Hummingbird.
- 346. AMAZILIA YUCATANENSIS (CABOT) GOULD.

  Buff-bellied Hummingbird.

1	88	PROCEEDINGS	OF UNITED	STATES	NATIONAL	MUSEUM.

- 347. BASILINNA XANTUSI (LAWR.) ELLIOT.

  Xantus's Hummingbird.
- 348. IACHE LATIROSTRIS (SWAINS.) ELLIOT.
  Broad-billed Hummingbird.
- 349. CYPSELUS SAXATILIS WOODH.
  White-throated Swift. [107.]
- 350. CYPSELOIDES NIGER BOREALIS (KENNERLY) RIDGW.
  Black Swift. [108.]
- 351. CHÆTURA PELAGICA (LINN.) BAIRD.
  Chimney Swift. [109.]
- 352. CHÆTURA VAUXII (Towns.) DE KAY.
  Vaux's Swift. [110.]
- 353. ANTROSTOMUS CAROLINENSIS (GM.) GOULD. Chuck-will's-widow. [111.]
- 354. CAPRIMULGUS VOCIFERUS (Wils.) Bp. Whip-poor-will. [112.]
- 355. PHALÆNOPTILUS NUTTALLI (Aud.) Ridgw.
  Poor-will. [113.]
- 356. NYCTIDROMUS ALBICOLLIS (GM.) BURM.

  Parauque Goatsucker. [116 a.]
- 357. CHORDEILES POPETUE (VIEILL.) Bb.
  Nighthawk. [114.]
- 357 a. CHORDEILES POPETUE HENRYI (Cass.) ALLEN. Western Nighthawk. [115.]
- 357 b. CHORDEILES POPETUE MINOR (CABAN.) RIDGW. Cuban Nighthawk.
- 358. CHORDEILES ACUTIPENNIS TEXENSIS (LAWR.) RIDGW.
  Texan Nighthawk. [116.]
- 359. CAMPEPHILUS PRINCIPALIS (LINN.) GRAY.

  Ivory-billed Woodpecker. [72.]
- 360. PICUS VILLOSUS LINN.

  Hairy Woodpecker. [74.]
- 360 a. PICUS VILLOSUS LEUCOMELAS (Bodd.) RIDGW.

  Great White-backed Sapsucker.
- 360 b. PICUS VILLOSUS HARRISI (AUD.) ALLEN.
  Harris's Woodpecker. [75.]
- 361. PICUS PUBESCENS LINN.
  Downy Woodpecker. [76.]
- 361a. PICUS PUBESCENS GAIRDNERI (Aud.) Coues. Gairdner's Woodpecker. [77.]

- 362. PICUS QUERULUS WILS.

  Red-cockaded Woodpecker. [80.]
- 363. PICUS SCALARIS WAGL.

  Texan Sapsucker. [79.]
- 363 a. PICUS SCALARIS LUCASANUS (XANT.) RIDGW. Saint Lucas Sapsucker.
- 364. PICUS NUTTALLI GAMB.

  Nuttall's Woodpecker. [78.]
- 365. PICUS STRICKLANDI MALH.
  Strickland's Woodpecker.
- 366. XENOPICUS ALBOLARVATUS (CASS.) BAIRD.
  White-headed Woodpecker. [51.]
- 367. PICOIDES ARCTICUS (SWAINS.) GRAY.

  Black-backed Three-toed Woodpecker. [82.]
- 368. PICOIDES TRIDACTYLUS AMERICANUS (BREHM) RIDGW.
  Banded-backed Three-toed Woodpecker. [83.]
- 368 a. PICOIDES TRIDACTYLUS DORSALIS (BAIRD) RIDGW.
  Striped-backed Three-toed Woodpecker. [84.]
- 369. SPHYRAPICUS VARIUS (LINN.) BAIRD.

  Yellow-bellied Woodpecker. [85.]
- 369a. SPHYRAPICUS VARIUS NUCHALIS BAIRD.

  Red-naped Woodpecker. [86.]
- 369 b. SPHYRAPICUS VARIUS RUBER (GM.) RIDGW.
  Red-breasted Woodpecker. [87.]
- 370. SPHYRAPICUS THYROIDEUS (Cass.) BAIRD.

  Black-breasted Woodpecker. [83, 89.]
- 371. HYLOTOMUS PILEATUS (LINN.) BAIRD.
  Pileated Woodpecker; Logcock. [90.]
- 372. CENTURUS CAROLINUS (L.) Bp.

  Red-bellied Woodpecker. [91.]
- 373. CENTURUS AURIFRONS WAGL.

  Golden-fronted Woodpecker. [92.7]
- 374. CENTURUS UROPYGIALIS BAIRD.
  Gila Woodpecker. [93.]
- 375. MELANERPES ERYTHROCEPHALUS (LINN.) Sw. Red-headed Woodpecker. [94.]
- 376. MELANERPES TORQUATUS (WILS.) BONAP.
  Lewis's Woodpecker. [96.]
- 377. MELANERPES FORMICIVORUS (Sw.) Bp. Californian Woodpecker. [95.]

- 190 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 377 a. MELANERPES FORMICIVORUS ANGUSTIFRONS BAIRD.
  Narrow-fronted Woodpecker.
- 378. COLAPTES AURATUS (LINN.) Sw.
  Yellow-shafted Flicker. [97.]
- 378 a. COLAPTES AURATUS HYBRIDUS (BAIRD) RIDGW. "Hybrid" Flicker. [98 a.]
- 378 b. COLAPTES AURATUS MEXICANUS (Sw.) RIDGW. Red-shafted Flicker. [98.]
- 379. COLAPTES CHRYSOIDES (MALH.) BAIRD.

  Malherbe's Flicker. [99.]
- 380. COLAPTES RUFIPILEUS RIDGW.
  Guadalupe Flicker.
- 381. MOMOTUS CÆRULEICEPS GOULD.

  Blue-capped Motmot. [119.]
- 382. CERYLE ALCYON (LINN.) BOIE.

  Belted Kingfisher. [117.]
- 383. CERYLE AMERICANA CABANISI (TSCHUDI) COUES.
  Texan Kingfisher. [118.]
- 384. TROGON AMBIGUUS GOULD.

  Coppery-tailed Trogon. [65.]
- 385. GEOCOCCYX CALIFORNIANUS (Less.) BAIRD.

  Road-runner; Chaparral Cock. [68.]
- 386. COCCYZUS SENICULUS (LATH.) VIEILL.

  Mangrove Cuckoo. [71.]
- 387. COCCYZUS AMERICANUS (LINN.) Br. Yellow-billed Cuckoo. [69.]
- 388. COCCYZUS ERYTHROPHTHALMUS (WILS.) BAIRD.
  Black-billed Cuckoo. [70.]
- 389. CROTOPHAGA ANI LINN.
  Savannah Blackbird. [66, 67.]
- 390. CROTOPHAGA SULCIROSTRIS SWAINS.
  Groove-billed Crotophaga.
- 391. RHYNCHOPSITTA PACHYRHYNCHA (SWAINS.) Bp.
  Thick-billed Parrot. [64.]
- 392. CONURUS CAROLINENSIS (LINN.) KUHL.

  Carolina Parakeet. [63.]
- 393. CONURUS HOLOCHLORUS BREVIPES BAIRD.
  Socorro Parakeet.
- 394. ALUCO FLAMMEUS AMERICANUS (AUD.) RIDGW.
  American Barn Owl. [47.]

- 395. ASIO AMERICANUS (STEPH.) SHARPE. American Long-eared Owl. [51.]
- 396. ASIO ACCIPITRINUS (PALL.) NEWTON. Short-eared Owl. [52.]
- 397. STRIX NEBULOSA FORST. Barred Owl. [54.]
- 397 a. STRIX NEBULOSA ALLENI RIDGW. Florida Barred Owl.
- 398. STRIX OCCIDENTALIS (XANT.) RIDGW. Spotted Owl.
- 399. ULULA CINEREA (GMEL.) Bp. Great Gray Owl. [53.]
- [399 a.] ULULA CINEREA LAPPONICA (RETZ.) RIDGW. Lapland Owl.
- 400. NYCTALE TENGMALMI RICHARDSONI (Bp.) RIDGW. Richardson's Owl. [55.]
- 401. NYCTALE ACADICA (GMEL.) Bp. Saw-whet Owl. [56, 57.]
- 402. SCOPS ASIO (LINN.) BP. Little Screech Owl. [49.]
- 402 a. SCOPS ASIO FLORIDANUS RIDGW. Florida Screech Owl.
- 402b. SCOPS ASIO MACCALLI (CASS.) RIDGW. Texan Screech Owl. [50.]
- 402 c. SCOPS ASIO MAXWELLIÆ RIDGW. Rocky Mountain Screech Owl.
- 402d. SCOPS ASIO KENNICOTTII (ELLIOT) RIDGW. Northwestern Screech Owl.
- 403. SCOPS TRICHOPSIS WAGL. Mexican Screech Owl.
- 404. SCOPS FLAMMEOLUS (LICHT.) Scl. Flammulated Screech Owl.
- 405. BUBO VIRGINIANUS (GM.) BP. Great Horned Owl. [48.]
- 405 a. BUBO VIRGINIANUS SUBARCTICUS (HOY) RIDGW. Western Horned Owl.
- 405 b. BUBO VIRGINIANUS ARCTICUS (SWAINS.) CASS. Arctic Horned Owl.
- 405 c. BUBO VIRGINIANUS SATURATUS RIDGW. Dusky Horned Owl.

- 192 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 406. NYCTEA SCANDIACA LINN.
  Snowy Owl. [61.]
- 407. SURNIA FUNEREA (LINN.) RICH & Sw. American Hawk Owl. [62.]
- [407 a.] SURNIA FUNEREA ULULA (LINN.) RIDGW.

  European Hawk Owl.
- 408. SPEOTYTO CUNICULARIA HYPOGÆA (BONAP.) RIDGW.
  Burrowing Owl. [58, 59.]
- 408a. SPEOTYTO CUNICULARIA FLORIDANA RIDGW. Florida Burrowing Owl.
- 409. GLAUCIDIUM GNOMA WAGL.

  California Pigmy Owl. [60.]
- 410. GLAUCIDIUM PHALÆNOIDES (DAUD.) SCL. & SALV. Ferruginous Pigmy Owl.
- 411. MICRATHENE WHITNEYI (COOPER) COUES.
  Whitney's Pigmy Owl.
- 412. HIEROFALCO GYRFALCO CANDICANS (GM.) RIDGW.
  White Gyrfalcon. [11.]
- 412a. HIEROFALCO GYRFALCO ISLANDUS (GM.) RIDGW. Iceland Gyrfalcon. [12.]
- 419 b. HIEROFALCO GYRFALCO SACER (FORST.) RIDGW. McFarlane's Gyrfalcon.
- 412 c. HIEROFALCO GYRFALCO OBSOLETUS (GM.) RIDGW. Labrador Gyrfalcon.
- 413. HIEROFALCO MEXICANUS POLYAGRUS (Cass.) Ridgw. Prairie Falcon. [10.]
- 414. FALCO PEREGRINUS NÆVIUS (GM.) RIDGW.

  American Peregrine Falcon; Duck Hawk. [5,6.]
- 414 a. FALCO PEREGRINUS PEALEI RIDGW.
  Peale's Falcon.
- 415. FALCO ALBIGULARIS DAUD.

  Chestnut-thighed Falcon. [8.]
- [416.] ÆSALON REGULUS (PALL.) BLYTH. European Merlin.
- 417. ÆSALON COLUMBARIUS (LINN.) KAUP.
  Pigeon Hawk. [7.]
- 417 a. ÆSALON COLUMBARIUS SUCKLEYI RIDGW. Black Merlin.
- 418. ÆSALON RICHARDSONII RIDGW.
  Richardson's Merlin.

- 419. RHYNCHOFALCO FUSCO-CÆRULESCENS (VIEILL.) RIDGW.
  Aplomado Falcon. [9.]
- 420. TINNUNCULUS SPARVERIUS (LINN.) VIEILL.
  Sparrow Hawk. [13.]
- 420 a. TINNUNCULUS SPARVERIUS ISABELLINUS (SWAINS.) RIDGW.
  Isabelline Sparrow Hawk.
- 421. TINNUNCULUS SPARVERIOIDES (VIG.) GRAY,
  Cuban Sparrow Hawk.
- [422.] TINNUNCULUS ALAUDARIUS (GM.) GRAY. European Kestril.
- 423. POLYBORUS CHERIWAY (JACQ.) CABAN.

  Caracara Eagle. [45.]
- 424. POLYBORUS LUTOSUS RIDGW.
  Guadalupe Caracara.
- 425. PANDION HALIAËTUS CAROLINENSIS (GM.) RIDGW.
  American Osprey; Fish Hawk. [44.]
- 426. **ELANOIDES FORFICATUS** (LINN.) RIDGW. Swallow-tailed Kite. [34.]
- 427. ELANUS GLAUCUS (BARTR.) COUES.
  White-tailed Kite. [35.]
- 428. ICTINIA SUBCÆRULEA (BARTR.) COUES.
  Mississippi Kite. [36.]
- 429. ROSTRHAMUS SOCIABILIS PLUMBEUS RIDGW. Everglade Kite. [37.]
- 430. CIRCUS HUDSONIUS (LINN.) VIEILL.

  Marsh Hawk. [38.]
- 431. ACCIPITER COOPERI BONAP.

  Cooper's Hawk. [15, 16.]
- 432. ACCIPITER FUSCUS (GMEL.) BP.
  Sharp-shinned Hawk. [17.]
- 433. ASTUR ATRICAPILLUS (Wils.) Bp.

  American Goshawk. [14.]
- 433 a. ASTUR ATRICAPILLUS STRIATULUS RIDGW.
  Western Goshawk.
- 434. ANTENOR UNICINCTUS HARRISI (Aud.) RIDGW.
  Harris's Hawk. [46.]
- [435.] BUTEO VULGARIS LEACH.
  European Buzzard.
- 436. BUTEO BOREALIS (GM.) VIEILL.

  Red-tailed Hawk. [23.]

  Proc. Nat. Mus. 80——13

Aug. 24, 1880.

- 194 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 436 a. BUTEO BOREALIS KRIDERI Hoopes. Krider's Hawk.
- 436 b. BUTEO BOREALIS CALURUS (Cass.) RIDGW.
  Western Red-tail. [20, 24.]
- 436 c. BUTEO BOREALIS LUCASANUS RIDGW. Saint Lucas Red-tail.
- 436 d. BUTEO BOREALIS SOCORROENSIS RIDGW. Socorro Red-tail.
- 437. BUTEO COOPERI Cass.

  Cooper's Henhawk. [29.]
- 438. BUTEO HARLANI Aud. Harlan's Hawk. [22.]
- 439. BUTEO LINEATUS (GM.) JARD.

  Red-shouldered Hawk. [25.]
- 439 a. BUTEO LINEATUS ELEGANS (CASS.) RIDGW.
  Red-bellied Hawk. [26.]
- 440. BUTEO ABBREVIATUS CABAN.

  Zone-tailed Hawk.
- 441. BUTEO ALBICAUDATUS VIEILL.
  White-tailed Hawk.
- 442. BUTEO SWAINSONI BONAP.

  Swainson's Hawk. [18, 19, 21, 28,]
- 443. BUTEO PENNSYLVANICUS (Wils.) Bp. Broad-winged Hawk. [27.]
- 444. URUBITINGA ANTHRACINA (LICHT.) LAFR.

  Mexican Black Hawk.
- 445. ASTURINA NITIDA PLAGIATA (LICHT.) RIDGW. Mexican Goshawk. [33.]
- 446. ONYCHOTES GRUBERI RIDGW.
  Gruber's Hawk.
- 447. ARCHIBUTEO LAGOPUS SANCTI-JOHANNIS (GMEL.) RIDGW.
  American Rough-legged Hawk. [30, 31.]
- 443. ARCHIBUTEO FERRUGINEUS (LICHT.) GRAY.

  Ferruginous Rough-leg. [32.]
- 449. AQUILA CHRYSAËTUS CANADENSIS (LINN.) RIDGW. Golden Eagle. [39.]
- 450. THRASAËTUS HARPYIA (LINN.) GRAY.
  Harpy Eagle.
- 451. HALIÆËTUS LEUCOCEPHALUS (LINN.) SAVIG. Bald Eagle; Gray Eagle. [41, 43.]

- 452. HALLÆËTUS ALBICILLA (LINN.) LEACH.
  Gray Sea Eagle. [42.]
- 453. PSEUDOGRYPHUS CALIFORNIANUS (SHAW) RIDGW.
  Californian Condor. [2.]
- 454. CATHARTES AURA (LINN.) ILLIG.
  Turkey Buzzard. [1.]
- 455. CATHARISTA ATRATA (Wils.) Less.

  Black Vulture; Carrion Crow. [3.]
- 456. COLUMBA PASCIATA SAY.

  Band-tailed Pigeon. [445.]
- 457. COLUMBA ERYTHRINA LIGHT.

  Red-billed Pigeon. [446.]
- 458. COLUMBA LEUCOCEPHALA LINN.
  White-crowned Pigeon. [447.]
- 459. ECTOPISTES MIGRATORIA (LINN.) Sw. Passenger Pigeon. [448.]
- 460. ZENÆDURA CAROLINENSIS (LINN.) Bp.
  Mourning Dove. [451.]
- 461. ZENÆDURA GRAYSONI BAIRD. Socorro Dove.
- 462. ZENÆDA AMABILIS Bp.

  Zenaida Dove. [449.]
- 463. ENGYPTILA ALBIFRONS (BP.) COUES.
  White-fronted Dove.
- 464. MELOPELIA LEUCOPTERA (L.) Bp.
  White-winged Dove. [450.]
- 465. CHAMÆPELIA PASSERINA (L.) SWAINS.
  Ground Dove. [453.]
- 466. SCARDAFELLA INCA (LESS.) BP. Scaled Dove. [452.]
- 467. GEOTRYGON MARTINICA (Gm.) Bp.
  Key West Dove. [454.]
- 468. STARNŒNAS CYANOCEPHALA (LINN.) Bp.
  Blue-headed Dove. [455.]
- 469. ORTALIS VETULA MACCALLI (BAIRD) RIDGW.
  Chachalaca; Texan Guan. [456.]
- 470. MELEAGRIS GALLOPAVO LINN.

  Mexican Turkey. [458.]
- 470 a. MELEAGRIS GALLOPAVO AMERICANA (BARTR.) COUES.
  Wild Turkey. [457.]

- 196 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 471. CANACE OBSCURA (SAY) Bp.

  Dusky Grouse. [459.]
- 471a. CANACE OBSCURA FULIGINOSA RIDGW. Sooty Grouse.
- 471 b. CANACE OBSCURA RICHARDSONII (Dougl.) BAIRD.
  Richardson's Grouse.
- 472. CANACE CANADENSIS (LINN.) Bp.

  Canada Grouse; Spruce Partridge. [460.]
- 472a. CANACE CANADENSIS FRANKLINI (DOUGL.) BAIRD. Franklin's Grouse. [461.]
- 473. BONASA UMBELLUS (LINN.) STEPH.
  Ruffed Grouse. [465.]
- 473 a. BONASA UMBELLUS UMBELLOIDES (DOUGL.) BAIRD.
  Gray Ruffed Grouse. [465 a.]
- 473 b. BONASA UMBELLUS SABINEI (DOUGL.) COUES.
  Oregon Ruffed Grouse. [466.]
- 474. LAGOPUS ALBUS (GM.) AUD.
  Willow Ptarmigan. [467, 470.]
- 475. LAGOPUS RUPESTRIS (GM.) LEACH.
  Rock Ptarmigan. [468.]
- 476. LAGOPUS LEUCURUS 8w.
  White-tailed Ptarmigan. [469.]
- 477. CUPIDONIA CUPIDO (LINN.) BAIRD. Prairie Hen. [464.]
- 477 a. CUPIDONIA CUPIDO PALLIDICINCTA RIDGW.
  Lesser Prairie Hen.
- 478. PEDIŒCETES PHASIANELLUS (L.) Elliot.

  Northern Sharp-tailed Grouse.
- 478 a. PEDIŒCETES PHASIANELLUS COLUMBIANUS (ORD) COUES.

  Common Sharp-tailed Grouse. [463.]
- 479. CENTROCERCUS UROPHASIANUS (Bp.) SWAINS.

  Sage Cock. [462.]
- 480. ORTYX VIRGINIANA (L.) Bp.

  Bob-white; American Quail. [471.]
- 480a. ORTYX VĮRGINIANA FLORIDANA COUES. Florida Quail.
- 480 b. ORTYX VIRGINIANA TEXANA (LAWR.) COUES.

  Texan Quail. [472.]
- 481. OREORTYX PICTA (Dougl.) BAIRD.

  Mountain Quail. [473.]

- 481 a. OREORTYX PICTA PLUMIFERA (GOULD) RIDGW.
  Plumed Quail.
- 482. LOPHORTYX CALIFORNICA (SHAW) BP.
  Californian Quail, [474.]
- 4S3. LOPHORTYX GAMBELI NUTT.

  Gambel's Quail. [475.]
- 484. CALLIPEPLA SQUAMATA (VIG.) GRAY. Scaled Quail. [476.]
- 485. CYRTONYX MASSENA (LESS.) GOULD.

  Massena Quail. [477.]
- 486. ARDEA OCCIDENTALIS AUD.

  Great White Heron; Würdemann's Heron. [488, 489.]
- 497. ARDEA HERODIAS LINN.

  Great Blue Heron. [487.]
- [488.] ARDEA CINEREA LINN.

  Common European Heron.
- 480. HERODIAS ALBA EGRETTA (GMEL.) RIDGW.
  American Egret. [486, 486 a.]
- 490. GARZETTA CANDIDISSIMA (GMEL.) Bp. Snowy Heron. [485.]
- 491. DICHROMANASSA RUFA (Bodd.) Ridgw.

  Reddish Egret; Peale's Egret. [482, 483.]
- 492. HYDRANASSA TRICOLOR LUDOVICIANA (WILS.) RIDGW.
  Louisiana Heron. [484.]
- 493. FLORIDA CÆRULEA (LINN.) BAIRD.
  Little Blue Heron. [490.]
- 494. BUTORIDES VIRESCENS (LINN.) Bp.
  Green Heron. [493.]
- 495. NYCTIARDEA GRISEA NÆVIA (BODD.) ALLEN. Black-crowned Night Heron. [495.]
- 496. NYCTHERODIUS VIOLACEUS (LINN.) REICH.
  White-crowned Night Heron. [496.]
- 497. BOTAURUS LENTIGINOSUS (MONTAG.) STEPH.
  American Bittern. [492.]
- 498. ARDETTA EXILIS (GMEL.) GRAY.
  Least Bittern. [491.]
- 499. MYCTERIA AMERICANA LINN.
  Jabiru.
- 500. TANTALUS LOCULATOR LINN.
  Wood Ibis. [497.]

- 198 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 501. EUDOCIMUS ALBUS (LINN.) WAGL.
  White Ibis. [499.]
- 502. EUDOCIMUS RUBER (LINN.) WAGL. Scarlet Ibis. [498.]
- 503. PLEGADIS FALCINELLUS (LINN.) KAUP.
  Glossy Ibis. [500.]
- 504. PLEGADIS GUARAUNA (LINN.) RIDGW.

  White-faced Glossy Ibis. [500 a.]
- 505. AJAJA ROSEA (Briss.) Ridgw.

  Roseate Spoonbill. [501.]
- [506.] HÆMATOPUS OSTRALEGUS LINN. European Oystercatcher.
- 507. HÆMATOPUS PALLIATUS TEMM.

  American Oystercatcher. [512.]
- 508. HÆMATOPUS NIGER PALL.

  Black Oystercatcher. [513.]
- 509. STREPSILAS INTERPRES (LINN.) ILLIG.
  Turnstone. [515.]
- 510. STREPSILAS MELANOCEPHALA VIG. Black Turnstone. [516.]
- 511. APHRIZA VIRGATA (GMEL.) GRAY.
  Surf Bird. [511.]
- [512.] VANELLUS CRISTATUS MEYER.
  Lapwing.
- 513. SQUATAROLA HELVETICA (LINN.) CUV Black-bellied Plover. [510.)
- [514.] CHARADRIUS PLUVIALIS LINN.
  Golden Plover.
- 515. CHARADRIUS DOMINICUS MÜLL.

  American Golden Plover. [503.]
- [515 a.] CHARADRIUS DOMINICUS FULVUS (GMEL.) RIDGW.
  Pacific Golden Plover.
- 516. OXYECHUS VOCIFERUS (LINN.) REICH.
  Killdeer. [504.]
- 517. ÆGIALITIS SEMIPALMATA (BONAP.) CABAN.
  Semipalmated Plover. [507.]
- 518. ÆGIALITIS HIATICULA (LINN.) BOIE. Ringed Plover.
- [519.] ÆGIALITIS CURONICA (GMEL.) GRAY.
  Little Ringed Plover.

- 520. ÆGIALITIS MELODA (ORD) BP.
  Piping Plover. [508.]
- 520a. ÆGIALITIS MELODA CIRCUMCINCTA RIDGW.
  Belted Piping Plover.
- 521. ÆGIALITIS CANTIANA NIVOSA (CASS.) RIDGW.
  Snowy Plover. [509.]
- 592. OCHTHODROMUS WILSONIUS (ORD) REICH.
  Wilson's Plover, [506.]
- 593. PODASOCYS MONTANA (Towns.) Cours.
  Mountain Plover. [505.]
- [524.] SCOLOPAX RUSTICULA LINN, European Woodcock,
- 525. PHILOHELA MINOR (GMEL.) GRAY,
  American Woodcock. [522.]
- [526.] GALLINAGO MEDIA LEACH.
  English Snipe.
- 526a. GALLINAGO MEDIA WILSONI (TEMM.) RIDGW. Wilson's Snipe. [523.]
- 527, MACRORHAMPHUS GRISEUS (GMEL.) LEACH.

  Red-breasted Snipe; Gray Snipe. [524.]
- 527 α. MACRORHAMPHUS GRISEUS SCOLOPACEUS (SAY) Coues.

  Red-bellied Snipe; Greater Gray-back. [525.]
- 528. MICROPALAMA HIMANTOPUS (BONAP.) BAIRD. Stilt Sandpiper. [536.]
- 529. TRINGA CANUTUS LINN.

  Knot; Robin Snipe. [526,]
- 530. ARQUATELLA MARITIMA (Brünn.) BAIRD.
  Purple Sandpiper. [528.]
- 531. ARQUATELLA COUESII RIDGW.
  Aleutian Sandpiper.
- 532. ARQUATELLA PTILOCNEMIS (Cours) Ridgw.
  Prybilov Sandpiper.
- [533.] ACTODROMAS ACUMINATA (Horsf.) Ridgw. Sharp-tailed Sandpiper.
- 534. ACTODROMAS MACULATA (VIEILL.) COUES.

  Pectoral Sandpiper. [531.]
- 535. ACTODROMAS COOPERI (BAIRD) COUES.

  Cooper's Sandpiper. [527.]
- 536. ACTODROMAS FUSCICOLLIS (VIEILL.) RIDGW Bonaparte's Sandpiper. [533.]

- 200 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 537. ACTODROMAS BAIRDII COUES.
  Baird's Sandpiper.
- 538. ACTODROMAS MINUTILLA (VIEILL.) Bp. Least Sandpiper. [532.]
- [539.] PELIDNA ALPINA (LINN.) BOIE. European Dunlin.
- 539 a. PELIDNA ALPINA AMERICANA CASS.

  Red-backed Sandpiper. [530.]
- [540.] PELIDNA SUBARQUATA (Guld.) Cuv. Curlew Sandpiper. [529.]
- 541. EREUNETES PUSILLUS (LINN.) CASS.

  Semipalmated Sandpiper. [535.]
- 541a. EREUNETES PUSILLUS OCCIDENTALIS (LAWR.) COUES.
  Western Sandpiper.
- 542. CALIDRIS ARENARIA (LINN.) ILLIG. Sanderling. [534.]
- 543. LIMOSA FEDOA (LINN.) ORD.

  Marbled Godwit. [547.]
- 544. LIMOSA LAPPONICA NOVÆ-ZEALANDIÆ GRAY.
  Pacific Godwit.
- 545. LIMOSA HÆMASTICA (LINN.) COUES.

  Hudsonian Godwit. [548.]
- [546.] LIMOSA ÆGOCEPHALA (LINN.) LEACH.

  Black-tailed Godwit.
- [547.] TOTANUS GLOTTIS (LINN.) BECHST.

  Green-shank. [538.]
- 548. TOTANUS MELANOLEUCUS (GMEL.) VIEILL.

  Greater Yellow-legs; Tell-tale. [539.]
- 549. TOTANUS FLAVIPES (GMEL.) VIEILL.
  Yellow-legs. [540.]
- 550. RHYACOPHILUS SOLITARIUS (WILS.) CASS.
  Solitary Sandpiper. [541.]
- [551.] RHYACOPHILUS OCHROPUS (LINN.) RIDGW.
  Green Sandpiper.
- 552. SYMPHEMIA SEMIPALMATA (GMEL.) HARTL. Willet. [537.]
- 553. HETEROSCELUS INCANUS (GMEL.) COUES.
  Wandering Tattler. [542.]
- [554.] MACHETES PUGNAX (Linn.) Cuv. Ruff. [544.]

- 555. BARTRAMIA LONGICAUDA (BECHST.) Bp.

  Bartram's Sandpiper; Field Plover. [545.]
- 556. TRYNGITES RUFESCENS (VIEILL.) CABAN.
  Buff-breasted Sandpiper. [546.]
- 557. TRINGOIDES MACULARIUS (LINN.) GRAY.
  Spotted Sandpiper. [543.]
- 558. NUMENIUS LONGIROSTRIS WILS.
  Long-billed Curlew. [549.]
- 559. NUMENIUS HUDSONICUS LATH.

  Hudsonian Curlew. [550.]
- 560. NUMENIUS BOREALIS (FORST.) LATH.
  Eskimo Curlew. [551.]
- [561.] NUMENIUS PHÆOPUS (LINN.) LATH.
  Whimbrel.
- [562.] NUMENIUS TAHITIENSIS (GMEL.) CASS.
  Bristle-thighed Curlew.
- 563. PHALAROPUS FULICARIUS (LINN.) Bp.
  Red Phalarope. [521.]
- 564. LOBIPES HYPERBOREUS (LINN.) Cuv.
  Northern Phalarope. [520.]
- 505. STEGANOPUS WILSONI (SAB.) COUES.
  Wilson's Phalarope. [519.]
- 566. RECURVIROSTRA AMERICANA GMEL.
  American Avocet. [517.]
- 567. HIMANTOPUS MEXICANUS (MÜLL.) ORD.

  Black-necked Stilt. [518.]
- 568. PARRA GYMNOSTOMA WAGL.

  Mexican Jacana.
- 569. RALLUS ELEGANS AUD.

  Red-breasted Rail. [542.]
- 570. RALLUS OBSOLETUS RIDGW.

  Californian Clapper Rail.
- 571. RALLUS LONGIROSTRIS CREPITANS (GMEL.) RIDGW.
  Clapper Rail. [553.]
- 571a. RALLUS LONGIROSTRIS SATURATUS HENSH.
  Louisiana Clapper Rail.
- 572. RALLUS VIRGINIANUS LINN.
  Virginian Rail. [554.]
- [573.] PORZANA MARUETTA (LEACH.) Bp. Spotted Crake.

- 202 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 574. PORZANA CAROLINA (LINN.) BAIRD.

  Sora Rail. [555.]
- 575. PORZANA NOVEBORACENSIS (GMEL.) BAIRD.
  Little Yellow Rail. [557.]
- 576. PORZANA JAMAICENSIS (GMEL.) BAIRD.
  Little Black Rail. [556.]
- 576a. PORZANA JAMAICENSIS COTURNICULUS BAIRD.
  Farallone Rail.
- [577.] CREX PRATENSIS BECHST.

  Corn Crake. [558.]
- 578. IONORNIS MARTINICA (LINN.) REICH.

  Purple Gallinule. [561.]
- 579. GALLINULA GALEATA (LICHT.) Bp. Florida Gallinule. [560.]
- 580. FULICA AMERICANA GMEL.

  American Coot. [559.]
- 581. ARAMUS PICTUS (BARTR.) COUES.

  The Limpkin. [481.]
- 582. GRUS AMERICANA (LINN.) TEMM.
  Whooping Crane. [478.]
- 583. GRUS CANADENSIS (LINN.) TEMM, Sandhill Crane. [479.]
- 584. GRUS FRATERCULUS CASS.
  Little Crane. [480.]
- 585. PHŒNICOPTERUS RUBER LINN.
  American Flamingo. [502.]
- [586.] OLOR CYGNUS (LINN.) Bp. European Swan.
- [587.] OLOR MINOR (PALL.) Bp.

  Bewick's Swan.
- 588. OLOR AMERICANUS (SHARPLESS) BP. Whistling Swan. [561 a.]
- 589. OLOR BUCCINATOR (RICH.) WAGL.
  Trumpeter Swan. [562.]
- 590. CHEN CÆRULESCENS (LINN.) RIDGW.
  Blue-winged Gcose. [564.]
- 591. CHEN HYPERBOREUS (PALL.) BOIE.

  Snow Goose. [563.]
- 591a. CHEN HYPERBOREUS ALBATUS (CASS.) RIDG.
  Lesser Snow Goose. [563a.]

- 592. CHEN ROSSII (BAIRD) RIDGW.
  Ross's Snow Goose.
- [593.] ANSER ALBIFRONS GMEL.

  European White-fronted Goose.
- 593 a. ANSER ALBIFRONS GAMBELI (HARTL.) COUES.

  American White-fronted Goose. [565, 566.]
- 594. BERNICLA CANADENSIS (LINN.) BOIE. Canada Goose. [567.]
- 594α. BERNICLA CANADENSIS HUTCHINSI (Sw. & Rich.) Woodh.
  Hutchins's Goose. [569.]
- 594b. BERNICLA CANADENSIS LEUCOPARIA (BRANDT) CASS.
  White-cheeked Goose. [568.]
- 594c. BERNICLA CANADENSIS OCCIDENTALIS (BAIRD) DALL & BANN.

  Larger White-cheeked Gcose. [567 a.]
- 595. BERNICLA BRENTA (PALL.) STEPH.
  Brant. [570.]
- 596. BERNICLA NIGRICANS (LAWR.) CASS.
  Black Brant. [571.]
- [597.] BERNICLA LEUCOPSIS (TEMM.) BOIL.
  Barnacle Goose. [572.]
- 598. PHILACTE CANAGICA (SEVAST.) BANNIST.
  Emperor Goose. [573.]
- 599. DENDROCYGNA AUTUMNALIS (Linn.) Evr.
  Black-bellied Tree Duck. [574.]
- 600. DENDROCYGNA FULVA (GMEL.) BURM.
  Fulvous Tree Duck. [575.]
- 601. ANAS BOSCAS LINN.

  Mallard. [576.]
- 602. ANAS OBSCURA GMEL.

  Black Duck. [577.]
- 603. ANAS FULVIGULA RIDGW.
  Florida Black Duck.
- 604. CHAULELASMUS STREPERUS (LINN.) GRAY.
  Gadwall. [584.]
- 605. DAFILA ACUTA (LINN.) BONAP.
  Pintail, [578.]
- [606.] MARECA PENELOPE (LINN.) SELBY.
  Widgeon. [586.]
  - 607. MARECA AMERICANA (GMEL.) STEPH.
    Baldpate. [585.]

- 204 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 608. SPATULA CLYPEATA (LINN.) BOIE. Shoveller. [583.]
- 609. QÜERQUEDULA DISCORS (LINN.) STEPH.
  Blue-winged Teal. [581.]
- 610. QUERQUEDULA CYANOPTERA (VIEILL.) CASS. Cinnamon Teal. [582.]
- [611.] NETTION CRECCA (LINN.) KAUP. English Teal. [580.]
  - 612. NETTION CAROLINENSIS (GMEL.) BAIRD.

    Green-winged Teal. [579.]
- 613. AIX SPONSA (LINN.) Boil.

  Wood Duck; Summer Duck. [587.]
- 614. FULIX MARILA (LINN.) BAIRD. Scaup Duck. [583.]
- 615. FULIX AFFINIS (Eyt.) BAIRD. Little Blackhead. [589.]
- 616. FULIX COLLARIS (Donov.) Baird.
  Ring-billed Blackhead. [590.]
- 617. AYTHYA VALLISNERIA (Wils.) Boie. Canvas-back. [592.]
- 618. AYTHYA AMERICANA (Evr.) Bp. Redhead. [591.]
- 619. CLANGULA ISLANDICA (GMEL.) Bp.

  Barrow's Golden-eye. [594.]
- 620. CLANGULA GLAUCIUM AMERICANA (Bp.) RIDGW.
  American Golden-eye. [593.]
- 691. CLANGULA ALBEOLA (LINN.) STEPH.

  Butterball; Bufflehead. [595.]
- 699. HISTRIONICUS MINUTUS (LINN.) DRESSER.
  Harlequin Duck. [596.]
- 623. HARELDA GLACIALIS (LINN.) LEACH.

  Long-tailed Duck; Old Squaw. [597.]
- 624. CAMPTOLÆMUS LABRADORIUS (GMEL.) GRAY.
  Labrador Duck. [600.]
- 625. POLYSTICTA STELLERI (PALL.) BRANDT. Steller's Duck. [598.]
- 626. LAMPRONETTA FISCHERI BRANDT. Fischer's Eider. [599.]
- 627. SOMATERIA MOLLISSIMA (LINN.) BOIE.

  Common Bider.

- 627 a. SOMATERIA MOLLISSIMA DRESSERI (SHARPE) COUES.
  American Eider. [606.]
- 628. SOMATERIA V-NIGRA GRAY.

  Pacific Eider. [607.]
- 629. SOMATERIA SPECTABILIS (LINN.) BOIE. King Eider. [608.]
- 630. ŒDEMIA AMERICANA Sw. & RICH.
  American Scoter. [604.]
- [631.] MELANETTA FUSCA (LINN.) BOIE.

  Velvet Scoter.
- 632. MELANETTA VELVETINA (CASS.) BAIRD.

  American Velvet Scoter. [601.]
- 633. **PELIONETTA PERSPICILLATA** (LINN.) KAUP. Surf Duck. [602.]
- 634. ERISMATURA RUBIDA (WILS.) Bp. Ruddy Duck. [609.]
- 635. NOMONYX DOMINICUS (LINN.) RIDGW.
  Black Masked Duck. [610.]
- 636. MERGUS MERGANSER AMERICANUS (Cass.) Ridgw.
  American Sheldrake. [611.]
- 637. MERGUS SERRATOR LINN.

  Red-breasted Sheldrake. [612.]
- 638. LOPHODYTES CUCULLATUS (LINN.] REICH.
  Hooded Sheldrake. [613.]
- 639. TACHYPETES AQUILA (LINN.) VIEILL.
  Frigate Pelican. [619.]
- 640. PELECANUS ERYTHRORHYNCHUS GMEL.
  American White Pelican. [615.]
- 641. PELECANUS FUSCUS LINN.

  Brown Pelican. [616.]
- 642. PHALACROCORAX CARBO (LINN.] Bp.

  Common Cormorant. [620.]
- 643. PHALACROCORAX DILOPHUS (Sw. & Rich.) NUTT.

  Double-crested Cormorant. [623.]
- 643a. PHALACROCORAX DILOPHUS FLORIDANUS (AUD.) RIDGW.
  Florida Cormorant. [624.]
- 643 b. PHALACROCORAX DILOPHUS CINCINNATUS (BRANDT) RIDGW.
  White-crested Cormorant. [622.]
- 644. PHALACROCORAX MEXICANUS (BRANDT) RIDGW.

  Mexican Cormorant. [625.]

206	PROCEEDINGS	of	UNITED	STATES	NATIONAL	MUSEUM.
-----	-------------	----	--------	--------	----------	---------

- 645. PHALACROCORAX PENICILLATUS (BRANDT) HEERM.
  Brandt's Cormorant. [626.]
- 646. PHALACROCORAX VIOLACEUS (GMEL.) RIDGW.
  Violet-green Cormorant. [627.]
- 646 a. PHALACROCORAX VIOLACEUS RESPLENDENS (Aud.) RIDGW.
  Baird's Cormorant.
- 647. PHALACROCORAX BICRISTATUS PALL.

  Red-faced Cormorant.
- 645. PHALACROCORAX PERSPICILLATUS PALL.
  Pallas's Cormorant. [621.]
- 649. PLOTUS ANHINGA LINN.
  American Anhinga; Snake Bird. [628.]
- 650. SULA BASSANA (LINN.) BRISS. Gannet. [617.]
- 651. SULA CYANOPS SUNDEY.

  Blue-faced Gannet.
- 652. SULA LEUCOGASTRA (Bodd.) Salv. Booby Gannet. [618.]
- 653. SULA PISCATOR (LINN.) Bp. Red-footed Booby.
- 654. PHAËTHON FLAVIROSTRIS BRANDT.
  Yellow-billed Tropic Bird. [629.]
- 655. PHAËTHON ÆTHEREUS LINN. Red-billed Tropic Bird.
- 656. RHYNCHOPS NIGRA LINN.

  Black Skimmer. [697.]
- 657. PAGOPHILA EBURNEA (PHIPPS) KAUP.

  Ivory Gull. [676, 677.]
- 658. RISSA TRIDACTYLA (LINN.) Bp. Kittiwake Gull. [672.]
- 658a. RISSA TRIDACTYLA KOTZBUEI (Bp.) Coues.
  Pacific Kittiwake.
- 650. RISSA BREVIROSTRIS BRANDT.

  Red-legged Kittiwake. [674, 675.]
- 660. LARUS GLAUCUS BRÜNN.
  Glaucous Gull; Burgomaster. [656.]
- 661. LARUS LEUCOPTERUS FABER.
  White-winged Gull. [658.]
- 669. LARUS GLAUCESCENS LICHT.

  Glaucous-winged Gull. [657,659.]

- 663. LARUS MARINUS LINN.

  Great Black-backed Gull. [600.]
- 664. LARUS OCCIDENTALIS AUD. Western Gull. [662.)
- [665.] LARUS AFFINIS REINH.
  Siberian Gull.
- 666. LARUS ARGENTATUS BRÜNN,
  Herring Gull.
- 666 a. LARUS ARGENTATUS SMITHSONIANUS COUES.

  American Herring Gull. [661.]
- 667. LARUS CACHINNANS PALL.
  Pallas's Herring Gull.
- 668. LARUS CALIFORNICUS LAWR.

  Californian Gull. [663.]
- 669. LARUS DELAWARENSIS ORD.

  Ring-billed Gull. [664.]
- 670. LARUS BRACHYRHYNCHUS RICH.
  Short-billed Gull. [664 a, 665, 673.]
- [671.] LARUS CANUS LINN.

  Mew Gull.
  - 672. LARUS HEERMANNI CASS.

    Heermann's Gull. [666.]
  - 673. LARUS ATRICILLA LINN.

    Laughing Gull. [667.]
- 674. LARUS FRANKLINI Sw. & RICH.
  Franklin's Gull. [668, 669.]
- 675. LARUS PHILADELPHIÆ (ORD) GRAY.

  Bonaparte's Gull. [670.]
- 676. RHODOSTETHIA ROSEA (MACGILL.) BRUCH.
  Ross's Gull. [678.]
- 677. XEMA SABINEI (J. SABINE) LEACH. Sabine's Gull. [680.]
- 678. CREAGRUS FURCATUS (NEB.) BP.
  Swallow-tailed Gull. [679.]
- 679. STERNA ANGLICA MONTAG.

  Gull-billed Tern. [681.]
- 680. STERNA CASPIA PALL.

  Caspian Tern. [682.]
- 681. STERNA REGIA GAMB.

  Royal Tern. [683.]

- 682. STERNA ELEGANS GAMB.

  Elegant Tern. [654.]
- 683. STERNA CANTIACA ACUFLAVIDA (CABOT) RIDGW.
  Cabot's Tern. [655.]
- 684. STERNA TRUDEAUI AUD.

  Trudeau's Tern. [687.]
- 685. STERNA FORSTERI NUTT.

  Forster's Tern. [691, 686. ]
- 686. STERNA FLUVIATILIS NAUM.
  Common Tern. [689.]
- 687. STERNA MACRURA NAUM.
  Arctic Tern. [690, 693.]
- 688. STERNA DOUGALLI MONTAG.
  Roseate Tern. [692.]
- 689. STERNA ALEUTICA BAIRD.
  Aleutian Tern.
- 690. **STERNA ANTILLARUM** (Less.) Coues. **Least Tern.** [694.]
- 691. STERNA FULIGINOSA GMEL. Sooty Tern. [688.]
- 692. STERNA ANÆSTHETA Scop. Bridled Tern.
- 693. HYDROCHELIDON LARIFORMIS SURINAMENSIS (GMEL.) RIDGW.
  Black Tern. [695.]
- [694.] HYDROCHELIDON LEUCOPTERA (Weisn. & Schinz) Boie. White-winged Black Tern.
- 695. ANOUS STOLIDUS LINN.

  Noddy Tern. [693.]
- 696. MEGALESTRIS SKUA (BRÜNN,) RIDGW. Skua Gull. [652]
- 697. STERCORARIUS POMATORHINUS (TEMM.) VIEILL Pomarine Jaeger. [653,]
- 698. STERCORARIUS PARASITICUS (LINN.) SCHEFF.
  Richardson's Jaeger. [654.]
- 699. STERCORARIUS BUFFONI (BOIE) COUES.
  Long-tailed Jaeger. [655.]
- 700. DIOMEDEA NIGRIPES AUD.

  Black-footed Albatross.
- 701. DIOMEDEA BRACHYURA TEMM.
  Short-tailed Albatross. [631.]

- [702.] DIOMEDEA CULMINATA GOULD.

  Yellow-nosed Albatross. [632.]
- 703. PHŒBETRIA FULIGINOSA (GMEL.) BP. Sooty Albatross. [633.]
- 704. OSSIFRAGA GIGANTEA (GM.) REICH.
  Giant Fulmar. [634.]
- 705. FULMARUS GLACIALIS (LINN.) STEPH, Fulmar Petrel. [635.]
- 705 a. FULMARUS GLACIALIS PACIFICUS (AUD.) BP.
  Pacific Fulmar. [636.]
- 705 b. FULMARUS GLACIALIS RODGERSI (CASS.) COUES.
  Rodger's Fulmar.
- 706. PRIOCELLA TENUIROSTRIS (Aub.) RIDGW.
  Slender-billed Fulmar. [637.]
- 707. PRIOFINUS MELANURUS (BONN.) RIDGW. Black-tailed Shearwater.
- [708.] PUFFINUS KUHLII (BOIE) BP.

  Cinereous Shearwater. [651.]
- 709. PUFFINUS MAJOR FABER.

  Greater Shearwater. [647.]
- 710. PUFFINUS CREATOPUS COOPER.
  Pink-footed Shearwater.
- [711.] PUFFINUS ANGLORUM TEMM.

  Manx Shearwater. [649.]
- 712. PUFFINUS AUDUBONII FINSCH.

  Dusky Shearwater. [650.]
- 713. PUFFINUS GAVIA (FORST.) FINSCH.

  Black-vented Shearwater.
- 714. PUFFINUS FULIGINOSUS STRICKL.
  Sooty Shearwater. [648.]
- 715. PUFFINUS GRISEUS (GM.) FINSCH.

  Dark-bodied Shearwater.
- 716. PUFFINUS TENUIROSTRIS TEMM.
  Slender-billed Shearwater.
- 717. ŒSTRELATA HÆSITATA (TEMM.) COUES.

  Black-capped Petrel. [638.]
- [718.] ŒSTRELATA BULWERI (JARD. & SELBY) COUES.
  Bulwer's Petrel.
- [719.] DAPTION CAPENSIS (LINN.) STEPH.

  Pintado Petrel; Cape Pigeon. [639.]

  Proc. Nat. Mus. 80——14 Sept. 4, 1880.

2	1	0	PROCEEDINGS	OF	UNITED	STATES	NATIONAL	MUSEUM

- 720. HALOCYPTENA MICROSOMA COUES.
  Least Petrel.
- 721. PROCELLARIA PELAGICA LINN.
  Stormy Petrel; Mother Carey's Chicken. [645.]
- 722. OCEANITES OCEANICA (KUHL) COUES.
  Wilson's Petrel. [644.]
- 723. CYMOCHOREA LEUCORRHOA (VIEILL.) COUES. Leach's Petrel. [642.]
- 724. CYMOCHOREA MELANIA (Bp.) Coues. Black Petrel. [643.]
- 725. CYMOCHOREA HOMOCHROA COUES.
  Ashy Petrel.
- 726. OCEANODROMA FURCATA (GMEL.) Bp. Fork-tailed Petrel. [640.]
- 727. OCEANODROMA HORNBYI (GRAY) Bp.

  Hornby's Petrel. [641.]
- 728. FREGETTA GRALLARIA (VIEILL.) Bp.
  White-bellied Petrel. [646.]
- 729. ÆCHMOPHORUS OCCIDENTALIS (LAWR.) COUES. Western Grebe. [704.]
- 730. ÆCHMOPHORUS CLARKII (LAWR.) COUES. Clark's Grebe. [705.]
- 731. PODICEPS HOLBÖLLI REINH.

  American Red-necked Grebe. [702,703a.]
- 732. DYTES CORNUTUS (GM.) KAUP. Horned Grebe. [706.]
- [733.] DYTES AURITUS (LINN.) RIDGW. Eared Grebe. [708.]
  - 733 a. DYTES AURITUS CALIFORNICUS (LAWR.) RIDGW.
    American Eared Grebe. [707.]
- 734. TACHYBAPTES DOMINICUS (LINN.) COUES. St. Domingo Grebe. [708a.]
- 735. PODILYMBUS PODICEPS (LINN.) LAWR.
  Thick-billed Grebe. [709.]
- 736. COLYMBUS TORQUATUS BRÜNN.
  Loon. [698.]
- 737.° COLYMBUS ADAMSI GRAY.

  Great White-billed Loon.
- 738. COLYMBUS ARCTICUS LINN.

  Black-throated Diver. [699.]

- 739. COLYMBUS PACIFICUS LAWR. Pacific Diver. [700.]
- 740. COLYMBUS SEPTENTRIONALIS LINN.
  Red-throated Diver. [701.]
- 741. ALCA IMPENNIS LINN.

  Great Auk. [710.]
- 742. UTAMANIA TORDA (LINN.) LEACH.
  Razor-billed Auk. [711.]
- 743. FRATERCULA ARCTICA (LINN.) STEPH.

  Common Puffin. [715, 716.]
- 743a. FRATERCULA ARCTICA GLACIALIS (LEACH) RIDGW.
  Large-billed Puffin. [714.]
- 744. FRATERCULA CORNICULATA (NAUM.) GRAY.
  Horned Puffin. [713.]
- 745. LUNDA CIRRHATA PALL.

  Tufted Puffin. [712.]
- 746. CERATORHINA MONOCERATA (PALL.) CASS.
  Horn-billed Puffin. [717,718.]
- 747. PHALERIS PSITTACULA (PALL.) TEMM.
  Parrot Auk. [725.]
- 748. SIMORHYNCHUS CRISTATELLUS (PALL.) MERREM.
  Crested Auk. [719, 720.]
- 749. SIMORHYNCHUS PYGMÆUS (GMEL.) RIDGW. Whiskered Auk. [721.]
- 750. CICERONIA PUSILLA (PALL.) RIDGW. Least Auk. [722, 723.]
- 751. PTYCORHAMPHUS ALEUTICUS (Pall.) Brandt.
  Cassin's Auk. [724.]
- 752. ALLE NIGRICANS LINK.

  Sea Dove; Dovekie. [738.]
- 753. SYNTHLIBORHAMPHUS ANTIQUUS (GM.) COUES.
  Black-throated Guillemot. [736.]
- 754. SYNTHLIBORHAMPHUS WURMIZUSUME (TEMM.) COUES.
  Temminck's Guillemot. [737.]
- 755. BRACHYRAMPHUS MARMORATUS (GM.) BRANDT.
  Marbled Guillemot. [732, 733.]
- 756. BRACHYRHAMPHUS KITTLITZI BRANDT.
  Kittlitz's Guillemot. [735.]
- 757. BRACHYRHAMPHUS HYPOLEUCUS XANTUS.

  Xantus's Guillemot.

- 212 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.
- 758. BRACHYRHAMPHUS CRAVERI (SALVAD.) COUES.
  Craver's Guillemot.
- 759. BRACHYRHAMPHUS BRACHYPTERUS BRANDT. Short-winged Guillemot. [734.]
- 760. URIA GRYLLE (LINN.) BRÜNN.
  Black Guillemot. [726.]
- 761. URIA COLUMBA (PALL.) CASS.

  Pigeon Guillemot. [727.]
- 762. URIA CARBO (PALL.) GRAY.
  Scoty Guillemot. [728.]
- 763. LOMVIA TROILE (Linn.) Brandt.

  Common Guillemot. [729, 730.]
- 763 a. LOMVIA TROILE CALIFORNICA (BRYANT) COUES.
  California Guillemot.
- 764. LOMVIA ARRA (PALL.) Bp.

  Thick-billed Guillemot.
- 764a. LOMVIA ARRA BRUNNICHI (Sch.) Ridgw. Brunnich's Guillemot. [731.]

# APPENDIX.

The following tables are intended as a condensed analysis of the changes which have taken place in North American ornithology since 1859, with other items of interest in the same connection.

a. Species eliminated from the catalogue of 1859.

- 4. Cathartes Burrovianus, Cassin Not North American?
- 6. Falco nigriceps, Cassin = No. 414.
- 16. ACCIPITER MEXICANUS, Swains. = No. 431.
- 19. Buteo bairdii, Hoy = No. 442, young.
- 21. Buteo insignatus, Cassin = No. 442, melanistic.
- 24. Buteo montanus, Nuttall = No. 436b, lighter phase.
- 28. Buteo oxypterus, Cassin = No. 442, young.
- 30. ARCHIBUTEO LAGOPUS, Gray = No. 447, light phase.
- 40. Haliaëtus pelagicus, Siebold. Not North American.
- 41. Haliaëtus washingtonii, Jard. = No. 451, young female.
- 56. NYCTALE ALBIFRONS, Cassin = No. 401, young.
- 59. ATHENE CUNICULARIA, Bon. The true cuniculariu is a South American form.\*
- 66. Crotophaga rugirostris, Sw. = No. 389.
- 73. Camperhilus imperialis, Gray. Not North American.
- 88. Sphyropicus Williamsonii, Baird = No. 370, adult male.
- 100. Lampornis Mango, Swains. Not North American.  $[=L.\ violicanda\ (Bodd.)\ Ell.]$
- 129. Tyrannus melancholicus, Vieill. Not North American.
- 167a. Var. MNIOTILTA LONGIROSTRIS, Baird.†
- 171. GEOTHLYPIS VELATUS, Cab. Not North American.
- 215. Myiodioctes bonapartei, Aud. = No. 127, young.
- 239. Collyrio elegans, Baird. An Asiatic species (Lanius lahtora, Sykes.).
- 242. Vireo virescens, Vieill. = No. 135?
- 253a. Var. Mimus caudatus, Baird. Not separable from polyglottus.
- 259a. Harporhynchus vetula, Baird = No. 15.
- 261a. HARPORHYNCHUS LONGICAUDA, Baird. Scarcely separable from rufus.
- 272. Troglodytes americanus, Aud. = No. 63.
- 289a. Var. Parus albescens, Baird = No. 41a.
- 309. Carpodacus haemorrhous, Wagl. Not North American?
- 311. Chrysomitris stanleyi, Bonap. Not North American. [= C. Barbata (Mol.).]
- 312. Chrysomitris Yarrelli, Bonap. Not North American.
- 324. Leucosticte arctous, Bonap. No sufficient evidence of occurrence in North America.
- 329. Plectrophanes melanomus, Baird = No. 189.
- 365. Melospiza Gouldii, Baird = No. 231c.
- 405. TRUPIALIS MILITARIS, Bonap. Not North American.

<sup>\*</sup>There appears to be but a single race inhabiting North America.

<sup>†</sup>While there is undoubtedly a very appreciable difference between specimens of *M. varia* from the West Indies and those from the *interior* of Eastern North America in the length of the bill, it is the small-billed form which should receive a new name, since Linnaus's name *varia* was based upon the bird of the South Atlantic States and West Indies. If to be regarded as separable, the name *borealis*, Nutt., may be applied to the western birds.

424. Corvus Cacalotl, Wagl. = No. 280.

470. LAGOPUS AMERICANUS, Aud. = No. 475.

482. Demiegretta Pealii, Baird = No. 491, white phase.

486a. Herodias egretta v. californica = No. 489.

488. ARDEA WURDEMANNII, Baird = No. 486, colored phase.

494. BUTORIDES BRUNNESCENS, Baird. Not North American.

514. Haematopus ater, Viellot. Not North American.

566. Anser frontalis, Baird = No. 593a, young.

603. Pelionetta trowbridgii, Baird = No. 633.

605. OIDEMIA BIMACULATA, Baird = No. 632, young.

630. DIOMEDEA EXULANS, Linn. Not North American.

659. Larus Chalcopterus, Lawr. = No. 662.

665. Larus Suckleyi, Lawr. = No. 670, young.

669. Chroicocephalus cucullatus, Br. = No. 674, young, second year.

671. Chroicocephalus minutus, Bruch. Not North American.

673. Rissa septentrionalis, Lawr. = No. 670, adult.

675. RISSA NIVEA, Bruch = No. 659.

677. Pagophila Brachytarsi, Hölb. = No. 657.

686. Sterna Havelli, Aud. = No. 685, winter plumage.

693. Sterna pikei, Lawr. = No. 687, young.

703. Podiceps Cristatus, Lath. Not North American?

703a. Podiceps cooperi, Lawr. = No. 731, young.

716. SAGMATORIHNA LABRADORIA, Cas. = No. 745, young.

718. CERORHINA SUCKLEYI, Cassin = No. 746, young.

720. Phaleris Tetracula, Stephens = No. 748, winter dress.

722. Phaleris Microceros, Brandt = No. 750, summer dress.

730. URIA RINGVIA, Brünnich = No. 763, individual phase.

733. Brachyrhamphus Wrangelli, Br. = No. 755, winter dress.

Seven of the above are included in Coues's "Check List of North American Birds" (1873), viz., numbers 88, 100, 309, 488, 563a, 603, 693, and 703; the equivalent numbers of the "Check List" being, respectively, 305, 274, 141a, 450, 480a, 518a, 568, and 609—some of them hearing a different name from that given in the Smithsonian catalogue. Besides the foregoing, there are given in the "Check List" the following untenable names:

146a. ÆGIOTHUS LINARIA (L.) Cab. var. fuscescens, Cs. = No. 179, midsummer dress.

157bis. Centronyx ochrocephalus, Aiken = No. 191, autumnal plumage.

[187.] Passer domesticus, Linn. An introduced species.

215a. Icterus spurius (L.) Bp. var. Affinis, Lawr. Not separable from I. spurius.

[283.] AGYRTRIA LINNEI (Bp.) —. Not North American. [= A. tobaci (Gm.) Ell.]

374a. Chamæpelia passerina (L.) Sw. rar. pallescens, (Bd.) Cs. Untenable race. 445ter [appendix]. Ibis thalassinus, Ridg. = No. 504, young.

b. Species and vaces described or added to the North American fauna since 1859.\*

6. Turdus iliacus, Linn.—Cf. Reinhardt, Ibis, 1861, 6. (Greenland; two examples.) 7a. Merula migratoria propinqua, Ridgw.—Turdus migratorious propinquas, Bull.

Nutt. Orn. Club, ii. Jan. 1877, 9. (Western U. S.)

 MERULA CONFINIS, Baird.—Turdus confinis, Review Am. B. i. 1864, 29. (Todos Santos, Cape St. Lucas.)

 HARPORHYNCHUS CINEREUS, Xantus.—Proc. Philad. Acad. 1859, 298. (Cape St. Lucas.)

14a. HARPORHYNCHUS CINEREUS BENDIREI, Coues.—Am. Nat. vii. June, 1873, 330, fig. 69. (Tucson, Arizona; C. Bendire.)

<sup>\*</sup>The new forms are in small capitals, the other additions in italics. In order to reduce the number of references to a minimum, only the original description, or the first North American record of a species is given. In some cases we have not been able to quote the first reference, but have done so whenever practicable.

215

- 15a. Harporhynchus curvirostris Palmeri, Ridgw.—H. curvirostris, var. palmeri, Ridgw. in Coues's Key, 1872, 351. (Arizona.)
- Harporhynchus Graysoni, Baird.—Cf. Lawr. Ann. Lyc. N. Y. x. Feb. 1871, —. (Socorro I.)
- Cyanecula succica (Linn.) Brehm.—Cf. Adams, Ibis, 1878, 422. (St. Michael's, Alaska; seven examples.)
- 31. Regulus obscurus, Ridgw.—*R. calendula obscurus*, Bull. U. S. Geol. & Geog. Surv. Terr. ii. No. 2, Apr. 1, 1876, 184. (Gnadalupe I., Lower California.)
- 33a. REGULUS SATRAPA OLIVACEUS, Baird.—R. satrapa, var. olicaccus, Baird, Review Am. Birds, i. July, 1864, 65 (in text). (Western United States.)
- 34. Phylloscopus borealis (Blas.) Dress.—Phyllopneuste kennicotti, Baird, Trans. Chicago Acad. i. 1869, 313, pl. 30, fig. 2. (St. Michael's, Alaska.)
- 44. Parus einetus, Bodd.—P. sibiricus (Gm.) Ridgw. Bull. Nutt. Orn. Club, ii. Jan. 1878, 37. (St. Michael's, Alaska; L. M. Turner.)
- 46a. Parus Rufescens Neglectus, Ridgw.—P. rufescens, β, neglectus, Proc. U. S. Nat. Mus. i. Apr. 25, 1879, 485. (Coast California.)
- CAMPYLORHYNCHUS AFFINIS, Xantus.—Proc. Philad. Acad. 1859, 298. (Cape St. Lucas.)
- 58a. Salpinctes obsoletus guadalupensis, Ridgw.—Bull. U. S. Geol. & Geog. Surv. Terr. ii. No. 2, Apr. 1, 1876, 185. (Guadalupe I., Lower California.)
- 59a. CATHERPES MEXICANUS CONSPERSUS, Ridgw.—C. mexicanus, var conspersus, Ridgw. Am. Nat. Oct. 1873, 602. (Middle Province of U. S.)
- 60b. Thryothorus ludovicianus miamensis, Ridgw.—*T. ludovicianus* (Lath.) var. miamensis, Am. Nat. ix. Aug. 1875, 469. (Miami River. E. Florida.)
- 61a. Thryomanes bewicki spilurus (Vig.) Baird.—Review Am. Birds, i. 1864, 126. (Pacific slope of United States.)
- 61b. Thryomanes bewicki leucogaster, Baird.—Review Am. B. i. 1864, 127. (Southern border of U. S.)
- 62. Thryomanes Brevicauda, Ridgw.—Bull. U. S. Geol. & Geog. Surv. Terr. ii. No. 2, Apr. 1, 1876, 186. (Guadalupe I., Lower California.)
- 64. Troglodytes insularis, Baird.—Cf. Lawr. Ann. Lyc. N. Y. x. Feb. 1871, 3, (Socotro I.)
- 65a. Anorthura troglodytes pacificus, Baird.—*T. hyemalis*, var. *pacificus*, Review Am. B. i. 1864, 145. (Pacific coast U. S.)
- ANORTHURA ALASCENSIS, Baird.—Troglodytes alascensis, Trans. Chicago Acad. i. 1869, 315, pl. 30, fig. 3. (St. George's Island, Alaska; W. H. Dall.)
- 67a. Telmatodytes palustris paludicola, Baird.—Cistothorus pulustris, var. paludicola, Review Am. B. i. 1864, 148. (Pacific coast U. S.)
- 69. Motacilla alba, Linn.—Cf. Reinhardt, Ibis, 1861, 6. (Greenland.)
- Budytes flava (Linn.) Gray.—Cf. Baird, Trans. Chicago Ac. i. 1869, 3, pl. 30, fig. 1. (St. Michael's, Alaska; Pease & Banuister.)
- Anthus pruteusis (Linn.) Beehst.—Cf. Paulsen, ed. Hölboll, Faun. Grönl. 1846,
   Reinh. Ibis, 1831, 6 (Greenland); B. B. & R. Hist. N. Am. B. i. 1874,
   (St. Michael's, Alaska.)
- 74a. Muiotilta varia borcalis (Nutt.) Ridgw. [See p. 213, foot-note.]
- 80. HELMINTHOPHAGA LAWRENCEI, Herrick. Proc. Philad. Acad. 1874, 220, pl. xv. (New Jersey.)
- 82. Helminthophaga leucobronchialis, Brewster.—Am. Sportsman, v. Oct., 1874; Bull. Nutt. Orn. Club, i. 1876, 1, plate. (Massachusetts.)
- 83. Helminthophaga lucie, Cooper.—Proc. Calif. Acad. Sci. July, 1861, 120. (Ft. Mojave, California.)
- 86a. Helminthophaga celata lutescens, Ridgw.—H. eelata, var. lutescens, Ridgw.
  Am. Johr. Sci. & Arts, third ser. iv. Dec. 1872, 457. (Pacific coast U. S.)
- Parula pitiayumi insularis (Lawr.) Ridgw.—Parula insularis, Lawr. Ann. Lyc. N. Y. x. Feb. 1871. (Socorro I., N. W. Mexico.)

- 89a. Parula Pitiayumi Nigrilora, Cones.—P. nigrilora, Bull. U. S. Geol. & Geog. Surv. Terr. iv. 1878, 11. (Hidalgo, Texas; G. B. Sennett.)
- Peucedramus olivaceus (Girand) Coues.—Sylvia olivacea, Girand, Texan Birds, 1841, 14, pl. vii. fig. 2. ("Texas.")—Dendraca olivacea, Hensh. Am. Sportsman, v. 328, Feb. 20, 1875; Orn. Wheeler's Exp. 1875, 202. (S. Arizona.)
- 103a. Dendræga dominica albilora, Baird.—D. Dominica, var. albilora, Am. Nat. vii. Oct. 1873, 605. (Mississippi Valley, south to Guatemala and Honduras.)
- 104. Dendræga gracle, Coues.—Cf. Baird, Review Am. B. i. 1864, 210. (Ft. Whipple, Arizona: Coues.)
- 106. DENDRŒCA CHRYSOPARIA, Sel. & Salv.—Cf. Baird, Review Am. B. i. 1864, 185, toot-note. (San Antonio, Texas; Heermann.)
- 113a. DENDRECA PALMARUM HYPOCHRYSEA, Ridgw.—Bull. Nutt. Orn. Club, Nov. 1876, 84, 85. (Atlantic States.)
- 116a. Siurus nævius notabilis, Grinnell.—Cf. Ridgw. Proc. U. S. Nat. Mus. iii. March 27, 1880, 12. (Black Hills, Wyoming.)
- 125a. WILSONIA PUSILLA PILEOLATA (Pall.) Ridgw.—Myiodioetes pusillus, var. pileolata, Ridgw. Am. Jour. Sci. & Arts, iv. Dec. 1872, 457; Am. Nat. vii. Oct. 1873, 607. (Pacific coast N. Am.)
- Cardellina rubrifrons (Giraud) Sel.—Muscicapa rubrifrons, Giraud. Texan Birds, 1841, pl. vii. fig. 1. ("Texas.")—Cardellina rubrifrons, Henshaw, Orn. Wheeler's Exp. 1875, 211. (Arizona.)
- \* 133. Basileuterus culicirorus (Licht.) Bp.—Muscicapa brasicri, Giraud, Texan Birds, 1841, pl. vi. fig. 2.
  - 134. Basileuterus bellii (Giraud) Scl.—Muscicapa belli, Giraud, Texan Birds, 1841, pl. iv. fig. 1.
  - 139a. Vireosylvia gilva swainsoni, Baird.—Vireo swainsoni, Baird, B. N. Am. 1858, 336, in text. (Pacific coast U. S.)
  - 141b. Lanivireo solitarius plumbeus (Cones) Allen.—Vircosylvia plumbea, Cones, Proc. Philad. Acad. 1866, 73. (Ft. Whipple, Arizona.)
  - 146. VIREO PUSILLUS, Coues.—Proc. Philad. Acad. 1866, 76. (Date Creek, Arizona.)
  - 147. VIREO VICINIOR, Coues.—Proc. Philad. Acad. 1866, 75. (Ft. Whipple, Arizona.)
  - 149b. Lanius Ludovicianus robustus, Baird.—Collurio Ludovicianus, var. robustus, Am. Nat. vii. Oct. 1873, 608. (California?)
  - 164a. Pyranga estiva cooperi, Ridgw.—Pyranga cooperi, Proc. Philad. Acad. 1869, 130. (S. W. United States.)
  - Pyrrhula Cassini, Baird.—P. coecinea, var. cassini, Trans. Chicago Acad. i. 1869,
     316, pl. 29, fig. 1. (Nulato, Alaska; W. H. Dall.)
  - 170a. Carpodaens frontalis rhodoeolpus (Caban.) Ridgw.—Cf. Ridgw. Am. Jour. Sci. & Arts, v. Jan. 1873, 39. (Coast of California.)
  - 171. Carpodacus amplus, Ridgw.—Bull. U. S. Geol. & Geog. Surv. Terr. ii. No. 2, Apr. 1, 1876, 187. (Guadalupe I., Lower Cal.; E. Palmer.)
  - 175a. LEUCOSTICTE TEPHROCOTIS LITTORALIS (Baird) Ridgw.—L. littoralis, Baird, Trans. Chicago Acad. i. 1869, 318, pl. 28, fig. 1. (Sitka, Alaska.)
  - Leucosticte atrata, Ridgw.—Am. Sportsman, July 18, 1874, 241. (Colorado; C. A. Aiken.)
  - 177. Leucosticte australis, Allen.—L. tephrocotis, var. australis, Allen. Cf. Ridgw. Bull. Essex Inst. v. Nov. 1873, 189. (Mt. Lincoln, Colorado.)
  - 178a. ÆGIOTHUS CANESCENS EXILIPES (Coues) Ridgw.—Egiothus cxilipes, Coues, Proc. Philad. Acad. Nov. 1861, 385. (Arctic America.)
  - 179a. Egiothus linaria holbölli (Brehm) Ridgw.—Linaria holbölli, Вкенм, Vög. Deutschl.
  - 180. Ægiothus Brewsteri, Ridgw.—Ægiothus (flavirostris var.) brewsterii, Ridgw. Am. Nat. July, 1872, 433. (Waltham, Mass.)
  - 182a. Astragalinus psaltria arizonæ (Coues) Ridgw.—C. mcxicana, var. arizonæ, Coues, Proc. Philad. Acad. 1866, 82. (Ft. Whipple, Arizona.)

- 192. Passerculus princeps. Maynard.—Am. Nat. vi. 1872, 637. (Ipswich, Mass.)
- 195. PASSERCULUS GUTTATUS, Lawr.—Am. Lyc. N. Y. viii. 1867, 473 (Cape St. Lucas).
  Cf. Cooper, Orn. Cal. i. 1870, 185.
- 197a. Poweetes gramineus confinis, Baird.—P. gramineus, var. confinis, Baird, B. N. Am. 1858, 448, in text.. (Western U. S.)
- 198a. Coturniculus passerinus perpallidus, Ridgw.—C. passerinus, var. perpallidus, Ridgw. in Coues's "Key", 1872, 137. (Western U. S.)
- 201a. Ammodromus caudacutus nelsoni, Allen.—Proc. Boston Soc. xvii. March, 1875, 93. (N. E. Illinois.)
- 203. Ammodromus nigrescens, Ridgw.—A. maritimus, var. nigrescens, Ridgw. Bull. Essex Inst. Dec. 1873, 198. (Indian R., Florida.)
- 204a. Chondestes grammiea strigatus (Sw.) Ridgw.
  - Mr. H. K. Coale, of Chicago, Ill., has lately called my attention to certain differences between eastern (typical) and western specimens of this species, which, upon examination of a large series, I find to be quite constant and sufficiently appreciable to warrant the recognition of a western race. Western birds being exactly like those from Mexico in those points in which they differ from eastern specimens, Swainson's name strigatus (Chondestes strigatus, Philos. Jour. i. 1827, 435), based upon the Mexican bird, is available for the western and southern race.
- 207a. Zonotrichia gambeli intermedia, Ridgw.—Z. leneophrys, var. intermedia, Ridgw. Bull. Essex Inst. Dec. 1873, 198. (Middle Province of U. S., north to Alaska.)
- 211a. SPIZELLA SOCIALIS ARIZONÆ, Coues.—S. socialis, var. arizonæ, Coues, Key, 1872, 143. (Ft. Whipple, Arizona.)
- 216. Junco Aikeni, Ridgw.—J. hyemalis, var. Aikeni, Ridgw. Am. Nat. Oct. 1873, 612, 614. (Mts. of Colorado; C. E. Aiken.)
- 219. Junco annectens, Baird.—Orn. Cal. i. 1870, 564. (Rocky Mts., Ft. Bridger to Arizona and New Mexico.)
- 223. Junco insularis, Ridgw.—Bull. U. S. Geol. & Geog. Surv. Terr. ii. No. 2, Apr. 1, 1876, 188. (Guadalnpe I., Lower Cal.; E. Palmer.)
- 225a. Amphispiza bellii nevadensis, Ridgw.—*Poospiza bellii*, var. *neradensis*, Ridgw. Bull. Essex Inst. Nov. 1873, 191. (Middle Province of U. S.)
- 226a. Peucæa æstivalis illinoensis, Ridgw.—P. illinoensis, Bull. Nutt. Orn. Club, Oct. 1879, 219. (Texas to S. Illinois.)
- 227. Peucæa arizonæ, Ridgw.—P. æstiralis, var. Arizonæ, Ridgw. Am. Nat. Oct. 1873, 615. (S. Arizona.)
- 229. Peucæa Carpalis, Coucs.—Am. Nat. vii. June, 1873, 322. (Tucson, Arizona; C. Bendire.)
- 230a. Peucæa Ruficeps Boucardi (Scl.) Ridgw.—*Cf.* Henshaw, Orn. Wheeler's Exp. 1874, 117. (S. Arizona and S. New Mexico.)
- 231d. Melospiza fasciata guttata (Nutt.) Ridgw.
  - This is the "M. rufina" of the old catalogue. The true M. rufina (Brandt) is a larger and darker form from Sitka, rediscovered since the publication of "Birds of North America" (1858).
- 231e. Melospiza fasciata rufina (Brandt) Ridgw.—Cf. B. B. & R. Hist. N. Am. B. ii. 1874, 29. (British Columbia to Sitka.)
- 232. Melospiza cinerea (Gmel.) Ridgw.—" M. insignis", BAIRD, Trans. Chicago Acad. i. 1869, 319, pl. 29, fig. 2. (Kadiak, Alaska; F. Bischoff.)
- 237a. Pipilo erythrophthalmus alleni, Coues.—P. alleni, Coues, Am. Nat. v. Aug. 1871, 363. (Florida.)
- 238e. Pipilo Maculatus consobrinus, Ridgw.—Bull. U. S. Geol. & Geog. Surv. Terr. ii. No. 2, Apr. 1, 1876, 189. (Guadalupe I., Lower Cal.)
- 238d. PIPILO MACULATUS CARMANI (Lawr.) Ridgw.—Pipilo carmani, LAWR. Ann. Lyc. N. Y. x. 1871. 7. (Socorto I., N. W. Mexico.)

- 240a. Pipilo fuscus albigula (Baird) Ridgw:—P. albigula, Baird, Proc. Philad. Acad. Nov. 1859, 305. (Cape St. Lucas.)
- 242a. CARDINALIS VIRGINIANUS IGNEUS, Baird.—C. igneus, Baird, Proc. Philad. Acad. Nov. 1859, 305. (Cape St. Lucas.)
- 253. Phonipara zena (Linn.) Bryant.—Cf. B. B. & R. Hist. N. Am. B. ii, 1874, 93. (Key West, Florida; H. W. Henshaw.)
- 258a. Molothrus ater obscurus (Gmel.) Cones.—M. obscurus, Cass. Proc. Philad. 1866, 18. (Lower California.)
- 259. Molothrus æneus (Wagl.) Cab.—Cf. MERRILL, Bull. Nutt. Orn. Club, i. July, 1876, 88. (Ft. Brown, Texas.)
- 263a. STURNELLA MAGNA MENICANA (Scl.) Ridgw.—Cf. Brewer, Bull. Nutt. Orn. Club, iii. July, 1878, 152. (Ft. Brown, Texas; J. C. Merrill.)
- 276. Quiscalus palustris (Swains.) Cassin.—'' Q. major", GAMBEL, Jour. Philad. Acad. i. 1847, 47. (Gulf of California.)\*
- 278b. QUISCALUS VERSICOLOR ÆNEUS, Ridgw.—Q. æneus, Ridgw. Proc. Philad. Acad. 1869, 134. (Mississippi Valley, Hudson's Bay Terr., Maine, etc.)
- [279.] Sturnus vulgaris, Linn.—Cf. Reinhardt, Ibis, 1861, 7. (Greenland.)
- 290a. Cyanocitta stelleri frontalis, Ridgw.—Cyanura stelleri, var. frontalis, Ridgw. Am. Jour. Sci. & Arts, third ser. v. Jan. 1873, 41, 43. (Sierra Nevada, California.)
- 290b. Cyanocitta stelleri annectens, Baird.—Cyanura stelleri, var. annectens, Baird, in B. B. & R. Hist. N. Am. Birds, ii. 1874, 281, in text. (Northern Rocky Mts.)
- 297a. Perisoreus canadensis capitalis, Baird.—*P. canadensis*, var. *capitalis*, Baird, Bull. Essex Inst. v. Nov. 1873, 193. (Rocky Mountains.)
- 297b. Perisoreus canadensis fumifrons, Ridgw.—Proc. U. S. Nat. Mus. iii. March 27, 1880, 5. (Coast of Alaska.)
- 298. Perisoreus obscurus, Ridgw.—*P. canadensis*, var. *obscurus*, Ridgw. Bull. Essex Inst. Nov. 1873, 194. (Northwest coast of U. S.)
- [299.] Alanda arvensis, Linn.—Cf. Dresser & Sharpe, Birds Eur. pt.—, and B. B. & R. Hist. N. Am. B. ii. 1874, 136. (Greenland and Bermuda.)
- 300a. Eremophila alpestris leucolæma, Coues.—Birds N. W. 1874, 38. (Interior plains N. Am.)
- 300b. Eremophila alpestris chrysolama (Wagl.) .— E. cornuta, var. chrysolama, BAIRD, B.N. Am. 1858, 403, in text. (Southwestern U. S.)
- 308. Pitangus derbianus (Kaup) Scl.—P. derbyanus, Coues, The Country, July 13, 1878, 184. (Lomita, Texas; G. B. Sennett.)
- Myiozetetes texensis (Giraud) Sel.—Muscieapa texensis, GIRAUD, Texan Birds, 1841,
   pl. 1. ("Texas.")
- 310. Myiodinastes luteiventris, Bonap.—Cf. Henshaw, Orn. Wheeler's Exp. 1875, 346, pl. xiv. (S. Arizona.)
- 319. Contopus pertinax, Cab. & Hein.—Cf. Coues, Proc. Philad. Acad. 1866, 60. (Ft. Whipple, Arizona.)
- 329. Empidonax fulvifrons (Giraud) Scl.—Muscicapa fulvifrons, GIRAUD, Texan Birds, 1841, pl. ii. ("Texas.")
- 329a. Empidonax fulvifrons pallescens (Coues) Ridgw.—Mitrephorus pallescens, Coues, Proc. Philad. Acad. 1863, 63. (F). Whipple, Arizona.)
- 331. Ornithiou imberbe (Scl.) Coues.—"O. incanescens", Coues, The Country, July 13, 1878, 184. (Lomita, Texas; G. B. Sennett.)
- 334. Eugenes fulgens (Sw.) Gould.—Cf. Henshaw, Am. Nat. Apr. 1874, 241; Orn. Wheeler's Exp. 1875, 379. (Mt. Graham, Arizona.)
- 341. Selasphorus alleni, Henshaw.—Bull. Nutt. Orn. Club, ii. 1877, 54. (Coast of California.)

<sup>\*</sup> The National Museum possesses an example of what is apparently this species from the coast of Louisiana.

- 342. Atthis heloisæ (Less. & Del.) Rich.—Cf. Elliot, Illustr. Am. B. i. 1869, xxi. xii. plate. (El Paso, Texas; J. H. Clarke.)
- 343. Stellula calliope, Gould.—Calothorax calliope, Xantus, Proc. Philad. Acad. 1859, 190. (Ft. Tejon, Cal.)
- 344. Calothorax lucifer (Sw.) Gray.—" Doricha enicura", HENSH. Am. Sportsman, v. 328, Feb. 20, 1875; Orn. Wheeler's Exp. 1875, 381. Cf. Lawr. Bull. Nutt. Orn. Club, ii. Oct. 1877, 108. (Camp Bowie, Arizona.)
- 345. Amazilia fuscicaudata (Fras.) Ridgw.—"Pyrrophæna riefferi", MERRILL, Bull. Nutt. Orn. Club, i. Oct. 1876, 88. (Ft. Brown, Texas.) (f. Ridgw. Proc. U. S. Nat. Mus. i. 1878, 147 (synonymy and diagnosis).
- 346. Amazilia yucatanensis (Cabot) Gould.—"A. eervineiventris", MERRILL, Bull. Nutt. Orn. Club, ii. Jan. 1877, 26. (Ft. Brown, Texas.) Cf. Ridgw. Proc. U. S. Nat. Mus. i. 1878, 148 (synonymy and diagnosis).
- 347. Basilinna xantusi (Lawr.) Elliot.—Amazilia xantusi, Lawr. Ann. N. Y. Lyc. vii. April, 1860, 109 (=  $\varphi$ ).—Heliopædica castancocanda, Lawr. t. c. 145 (=  $\vartheta$ ). (Cape St. Lucas.)
- 348. Iaehe latirostris (Sw.) Elliot.—Ciree latirostris, Hensh. Am. Sportsman, v. Feb. 20, 1875; Orn. Wheeler's Exp. 1875, 380. (Chiracahua Mts., S. Arizona.)
- 357b. Chordeiles popetue minor (Cab.) Ridgw.—(f. B. B & R. Hist. N. Am. B. iii. 1874, 520. (Miami, Florida; C. J. Maynard.)
- 360a. Pieus villosus leucomelas (Bodd.) Ridgw.—Pieus leucomelas, Bodd. Tabl. P. E. 1783 (ex. Pl. Enlum. 345, fig. 1 = ♀ ad.).
- 363a. PICUS SCALARIS LUCASANUS (Xant.) Cones.—P. lucasanus, Xantus, Proc. Philad. Acad. 1859, 298, 302. (Cape. St. Lucas.)
- 365. Pieus stricklandi, Malh.—Cf. Henshaw, Am. Sportsman, v. 328, Feb. 20, 1875; Orn. Wheeler's Exp. 1875, 389. (S. Arizona.)
- 377a. Melanerpes formicivorus angustifrons, Baird.—M. farmicivorus, var. angustifrons, Baird, Orn. Cal. i. 1870, 405. (Cape St. Lucas.)
- 380. Colaptes rufipileus, Ridgw.—C. mexicanus rufipileus, Bull. Geog. & Geol. Surv. Terr. ii. No. 2, Apr. 1, 1876, 191. (Guadalupe I., Lower Cal.)
- 390. Crotophaga sulcirostris, Swains.—Cf. Coues, The Country, July 13, 1878, 184. (Lomita, Texas; G. B. Sennett.)
- 393. CONURUS HOLOCHLORUS BREVIPES, Baird.—Conurus holochlorus, var. brevipes, "Baird, MS.", LAWR. Ann. Lyc. N. Y. x. 1871, —. (Socorro I.)
- 397a. STRIX NEBULOSA ALLENI, Ridgw.—Proc. U. S. Nat. Mus. iii. March 27, 1880, —. (Clearwater, S. Florida.)
- 398. STRIX OCCIDENTALIS (Xant.) Ridgw.—Syrnium occidentale, Xantus, Proc. Philad. Acad. 1859, 193. (Ft. Tejon, Cal.)
- 399a. Ulula einerea lapponica (Retz.) Ridgw.—Syrnium lapponicum, Ridgw. Bull. Nutt. Orn. Club, iii. Jan. 1878, 37. (St. Michael's, Alaska; L. M. Turner.)
- 402a. Scops asio floridanus, Ridgw.—S. asio, var. Floridanus, Ridgw. Bull. Essex Inst. Dec. 1873, 200. (Florida.)
- 402c. Scops asio Maxwellle, Ridgw.—S. asio, ε, maxwelliw, Ridgw. Field & Forest, June, 1877, 210, 213. (Boulder Co., Colorado.)
- 402d. Scors asio Kennicottii, (Elliot) Coues.—S. Kennicottii, Elliot, Proc. Philad. Acad. 1867, 69; Illustr. Am. B. 1869, pl. 11. (Sitka, Alaska; F. Bischoff.)
- 403. Scops trichopsis, Wagl.—"S. asio, var. maecalli", B. B. & R. Hist. N. Am. B. iii. 1874, 52. (New Mexico.)
- 404. Scops flammeolus (Licht.) Scl.—Cf. Cooper, Orn. Cal. i. 1870, 422. (Ft. Crook, N. California.)
- 405a. Bubo virginianus subarcticus (Hoy) Ridgw.—Bubo subarcticus, Hoy, Proc. Philad.

  Acad. vi. 1852, 211. (Wisconsin.) [="var. arcticus" of Hist. N. Am. B. iii.
  1874, 64.]
- 405b. Bubo virginianus arcticus (Swains.) Cass.—Bubo arcticus, Swains. F. B. A. ii. 1831, 86, pl. 30. (Interior of fur countries.)

- 405c. Bubo virginianus saturatus, Ridgw.—Orn. 40th Parallel, 1877, 572, foot-note.

  (Northern coast N. Am.) [="var. pacificus" of Hist. N. Am. B. iii. 65.]
- 407a. Surnia funcrea ulula (Linn.) Ridgw.—Cf. Ridgw. Bull. Nutt. Orn. Club, iii. Jan. 1878, 38. (St. Michael's, Alaska; L. M. Turner.)
- 408a. Spectyto cunicularia floridana, Ridgw.—S. cunicularia, var. floridana, Ridgw. Am. Sportsman, iv. No. 14, July 4, 1874, 216. (Sarasota Bay, Florida.)
- 410. Glaucidium phalwooides (Daud.) Sel. & Salv.—G. ferrugineum (Max.) Coues, Am. Nat. vi. 1872, 370. (Tueson, Arizona; C: Bendire.)
- 411. MICRATHENE WHITNEYI (Cooper) Cones.—Athene whitneyi, Cooper, Proc. Cal. Acad. Sci. ii. 1863, 118. (Ft. Mojave, S. E. California.)
- 412b. Hierofalco gyrfalco sacer (Forst.) Ridgw.—F. sacer, Forster, Philos. Trans. lxii. 1772, 383, 423. (Hudson's Bay Terr.)
- 412c. Hierofaleo sacer obsoletus (Gm.) Ridgw.—Falco obsoletus, Gmel. S. N. i. 1788, 268. (Hudson's Bay Terr.)
- 414a. Falco peregrinus pealei, Ridgw.—F. communis, var. Pealei, Ridgw. Bull. Essex Inst. Dec. 1873, 201. (Northwest coast N. Am.)
- 416. Æsalon regulus (Pall.) Blyth.—Falco æsalon Newton, Man. Nat. Hist. Greenl. 1875, p. 96. (At sea, near Greenland, lat. 57° 41′ N., long. 35° 23′ W.)
- 417a. ÆSALON COLUMBARIUS SUCKLEYI, Ridgw.—Falco columbarius, var. Suckleyi, Ridgw. Bull. Essex Inst. v. Dec. 1873, 201. (Northwest coast N. Am.)
- 418. ÆSALON RICHARDSONH, Ridgw.—Falco (Hypotriorchis) richardsonii, RIDGW. Proc. Philad. Acad. Dec. 1870, 145. (Interior of North America.)
- 421. Tinnunculus sparverioides (Vig.) Gray. (Florida.)\*
- 422. Tinnunculus alaudarius (Gm.) Gray.—Cf. Newton, Man. Nat. Hist. Greenl. 1875, 96. (Off Cape Farewell, Greenland.)
- 424. POLYBORUS LUTOSUS, Ridgw.—Bull. U. S. Geol. & Geog. Surv. Terr. No. 6, 2d ser. Feb. 8, 1876, 459. (Guadalupe I., Lower California.)
- 433a. Astur atricapillus striatulus, Ridgw.—A. palumbarius, var. striatulus, Ridgw. in Hist. N. Am. B. iii. 1874, 240. (Western N. Am.)
- 435. Buteo rulgaris, Leach.—Cf. MAYNARD, Bull. Nutt. Orn. Club, i. No. 1, April, 1876, 2-6. (Pawpaw, Mich.)
- 436a. Buteo Borealis Krideri.—"B. borealis, variety kriderii," Hoopes, Proc. Philad. Acad. 1873, 238, pl. v. (=juv.; Winnebago Co., Iowa.)
- 436c. Buteo Borealis Lucasanus, Ridgw.—*B. borealis*, var. *lucasanus*, Hist. N. Am. B. iii. 1874, 258, 285. (Cape St. Lucas.)
- 436d. Buteo borealis socorroensis.

In the "Proceedings" of the Boston Society of Natural History, 1871, p. 42, Mr. Lawrence refers to a "Buteo borcalis, var. montanus, Nntt." as being very abundant on the island of Socorro, where it is the only species of hawk to be found, and where it is a "constant resident, rearing its young, and subsisting entirely on land crabs", etc. In the same paper, p. 10. Mr. Lawrence applies the same name to a lawk occurring abundantly on the Tres Marias Islands, and there subsisting "entirely upon the Iguana lizard and rabbits." In "History of North American Birds", vol. iii, p. 285 (1874), I referred the Tres Marias bird to Buteo borealis var costaricensis, and described our only specimen from that locality as the young of that form, although I had not, from want of specimens, been able to compare it with the corresponding age of the Central American bird. I now have strong doubts as to its being the same as B. costaricensis, while as to the hawk found on Socorro I regard it quite certain that it is, like the Caracara of Guadalupe (Polyborus lutosus), a species or race peculiar to that remote island, the birds of which are for the most part entirely local. I therefore propose to name, provisionally, the Socorro Hawk B. borealis socorroensis.

440. Buteo abbreviatus, Cabanis, in Schomb. Guiana, iii. 1848, 730.—"B. zonocereus, Sel.", Cooper, Orn. Cal. i. 1870, 479. (Coast of California, near San Diego.)

<sup>\*</sup>There is a male of this species in my collection said to have been obtained in Florida. It was found with a collection of Florida birds, forming part of a dealer's stock, and was evidently the same "make" of skin as the rest of the lot.

441. Butco albicaudatus (Vieill.)—B. albocaudatus, Cours, The Country, i. 184, July 13, 1878. (Lomita, Texas; G. B. Sennett). Cf. Ridgw. Proc. U. S. Nat., Mus. i. Oct. 2, 1878, 154 (synonymy and descriptions).

444. Urubitinga anthracina (Licht.) Nitzsch.—Cf. Henshaw, Am. Sportsman. v. 328. Feb. 20, 1875; Orn. Wheeler's Exp. 1875, 420. (Arizona.)

446. Onychotes Gruberi, Ridgw.—Proc. Philad. Acad. Dec. 1870, 149. (California.)

450. Thrasætus harpyia.

According to Dr. Felix L. Oswald, in the American Naturalist, 1878, p. 151, a specimen of the Harpy Eagle was shot at the "delta of the Rio Grande", in Texas, by Professor S. B. Buckley, State geologist of Texas. A full account of the circumstance is given in Dr. Oswald's interesting article. I have seen somewhere a record of the occurrence of this species in Louisiana, but cannot now lay hand on the reference. According to my recollection, the record may be found in an old number of the "Proceedings" of the Philadelphia Academy of Sciences, or else of the Zoölogical Society of London.

- 461. ZENÆDURA GRAYSONI, Baird.—Cf. LAWRENCE, Ann. Lyc. N. Y. x. 1871, 17; Proc. Boston Soc. xiv. 1871, 299. (Socorro I.)
- 463. Engyptila albifrons (Bp.) Coues.— Echmoptila albifrons, Coues, Bull. U. S. Geol. & Geog. Surv. Terr. iv. No. 1, 1878, 48 (South Texas); Ridgw. Proc. U. S. Nat. Mus. i. Oct. 1878, 158 (synonymy).
- 471a. Canace obscura fuliginosa, Ridgw.—C. obscura, var. fuliginosa, Ridgw. Bull. Essex Inst. Dec. 1873, 199. (Northwest coast, Oregon to Sitka.)
- 471b. Canace obscura richardsonii (Dougl.) Ridgw.—Tetrao richardsonii, "Sabine, MSS.", Dougl. Linn. Trans. xvi. 1829, 141.
- 477a. Cupidonia cupido pallidicineta, Ridgw.—C. cupido, var. pallidicineta, Ridgw. Bull. Essex Inst. Dcc. 1873, 199. (Southwestern prairies.)
- 478a. Pediocetes phasianellus columbianus (Ord) Ridgw.—Phasianus columbianus, Ord, Guthrie's Geog. 2d Am. ed. ii. 1815, 317.—Pediacetes columbianus, Elliot, Pr. Ac. Nat. Sci. Philad. 1862, 403.
- 480a. Ortyx virginiana floridana, Coues.—O. virginianus, var. floridanus, Coues, Key, 1872, 237. (Florida; J. A. Allen.)
- 481a. Oreortyx picta plumifera (Gould) Ridgw.—Ortyx plumifera, Gould, P. Z. S. 1837, 42.
- 488. Ardea cinerea, Linn.—Cf. Reinii. Ibis, 1861, 9. (Greenland.)
- 499. Mycteria americana, Linn.—Cf. Coues, Check List, 1873, 135. (Austin, Texas.)
- 506. Hamatopus ostralegus, Linn.—Cf. Reinh. Ibis, 1861, 9. (Greenland.)
- 512. Vanellus cristatus (Linn.) Meyer.—Cf. Reinh. Ibis, 1861, 9. (Greenland.)
- 514. Charadrius pluvialis, Linn.—Cf. Newton, Man. Nat. Hist. Greenl. 1875, 101.
- 515a. Charadrius dominicus fulrus (Gmel.) Ridgw.—C. fulrus, Coues, Elliott's Prybilov Islands, 1875, 179; Birds N. W. 1874, 450, foot-note. (Prybilov Islands, Alaska.)
- 518. Ægialitis hiaticula (Linn.) Boiè.—Cf. NEWTON, Man. Nat. Hist. Greenl. 1875, 101. (Greenland.)
- 519. Agialitis curonica (Gmel.) Gray.—Introduced as A. microrhynchus, Ridgw., n. s., Am. Nat. viii. Feb. 1874, 109. ("San Francisco, Cal.")
- 520a. ÆGIALITIS MELODA CIRCUMCINCTA, Ridgw.—.E. melodus, var. circumcinctus, Am. Nat. viii. Feb. 1874, 109. ("Plains between Missouri River and Rocky Mountains.")
- 524. Scolopax rusticula, Linn.—Cf. BAIRD, Am. Jour. Arts & Sciences, xli. May, 1866, 25. (Newfoundland.)
- 526. Gallinago media, Leach.—Cf. Reinh. Ibis, 1861, 11. (Greenland.)
- 531. ARQUATELLA COUESH, Ridgw.—Bull. Nutt. Orn. Club, July, 1880, 160. (Aleutian islands and contiguous coast of Alaska.)
- 532. Arquatella Ptilocnemis (Coues) Ridgw.—"Tringa crassirostris", Dall, Am. Nat. viii. 1873, 635 (St. Paul's I., Alaska).—Tringa ptilocnemis, Coues, Elliott's Prybilov Islands, 1875, foot-note.

533. Actodromas acuminata (Horsf.) Ridgw.—Obtained at St. Michael's, Alaska, by Mr. E. W. Nelson, U. S. Signal Service.

The single example forwarded by Mr. Nelson is without date or exact locality, and I have been unable to get from him the desired information, my letter to him on the subject having doubtless miscarded. The species is so important an addition to the fauna, however, that it cannot be overlooked in the present connection, reluctant as we are to publish a note of its occurrence without being able to furnish those data from the pen of Mr. Nelson himself, which would do him full justice, as the first discoverer of this Indian species on this continent.

- 537. ACTODROMAS BAIRDII, Cones.—Proc. Philad. Acad. 1861, 194. (Arctic America.)
- Pelidna alpina (Linn.) Boie. Tringa alpina, Newton, Man. Nat. Hist. Greenl. 1875, 103. (Greenland.)
   EREUNETES PUSILLUS OCCIDENTALIS (Lawr.) Cones. — E. occidentalis, Lawr. Proc.
- Philad. Acad. 1864, 107. (Pacific coast U. S.)
- 544. Limosa lapponica nova-zealandia, Gray.—L. uropygialis, BAIRD, Trans. Chicago Acad. i. 1869, 320, pl. 32. (Alaska.)
- 546. Limosa agocephala (Linn.) Leach.—Cf. REINH. Ibis, 1861, 11. (Greenland.)
- 551. Rhyacophilus ochropus (Linn.) Ridgw.—Cf. Brewer, Bull. Nutt. Orn. Club, iii. Jan. 1878, 49. (Nova Scotia.)
- 561. Numenius phaopus (Linn.) Lath.—Cf. Reinii. Ibis, 1861, 10. (Greenland.)
- 562. Numenius tahitiensis (Gmel.) Cass.—"Numenius femoralis, Peale", Ridgw. Am. Nat. July, 1874, 435. (Kadiak, Alaska; F. Bischoff.)
- 568. Parra gymnostoma, Wagl.—Cf. MERRILL, Bull. Nutt. Orn. Club, i. Nov. 1876, 88 (Ft. Brown, Texas); Ridgw. Proc. U. S. Nat. Mus. i. 1878, 167, pl. iii. (synonymy and descriptions).
- 570. Rallus obsoletus, Ridgw.—R. elegans, var. obsoletus, Ridgw. Am. Nat. viii. Feb. 1874, 111. (Coast California.)
- 571a. Rallus longirostris saturatus, Hensh.—Cf. Ridgw. Bull. Nutt. Orn. Club, July, 1880, 140. (Louisiana.)
- 573. Porzana maruetta (Leach) Bp.—"Ortygometra porzana (Linn.)", REINH. Ibis, 1861, 12. (Greenland.)
- 576a. Porzana Jamaicensis coturniculus, Baird.—P. jamaicensis, var. coturniculus,
   Baird, Am. Nat. viii. Feb. 1874, 111. (Farallone Islands, California.)
- 586. Olor cygnus (Linn.) Bp.—"Cygnus ferus, RAY", REINH. Ibis, 1861, 13. (Greenland.)
- 587. Olor minor (Pall.) Bp.—"Cygnus bewickii", Sw. & Rich. F. B. A. ii. 1831, 465. ("Igloolik [Arctic America], lat. 66°, June 19, 1823." Said to breed "on the sea-coast within the Arctic circle." The description, from specimens killed at locality quoted above, is of the true O. minor, or Bewick's Swan.)
- 592. CHEN ROSSII (Baird) Ridgw.—Anser rossii, "BAIRD, MSS.", CASS. Proc. Philad. Acad. 1861, 73. (Arctic America.)
- 593. Anser albifrons (Gm.) ——.—Cf. REINH. Ibis, 1861, 12. (Greenland.)
- 603. Anas fulvigula, Ridgw.—A. obscura, var. fulvigula, Am. Nat. viii. Feb. 1874, 111. (Florida.)
- 627a. Somateria mollissima dresseri (Sharpe) Cones.—S. dresseri, Sharpe, Ann. Mag. N. H. July, 1871, 51, figs. 1, 2.

This bird has been called by all American writers *S. mollissima*. The true Eider, however, has only lately been detected in America, it being the form found by Mr. Kumlien breeding abundantly on the west side of Cumberland Gulf.

- 631. Melanetta fusca (Linn.)—Cf. REINHARDT, Vid. Medd. Nat. För. Kjobenhavn, 1879, 1. (South Greenland.)
- 646a. Phalacrocorax riolaceus resplendens (And.) Ridgw.—Graculus bairdii, "Gryber, MSS.", Cooper, Proc. Philad. Acad. Jan. 1865, 5. (Farallone Islands.)

- 647. Phalaerocorax bicristatus, Pallas.—"Graenlus bicristatus, Pallas", Baird, Trans. Chicago Acad. i. 1869, —, pl. 33. (St. George's I., Alaska; W. H. Dall.)
- 651. Sula cyanops, Sundev.—Cf. Lawr. Proc. Boston Soc. xiv. 1871, 302. (Socorro I.)
- 653. Sula piseator (Linu.) .- Cf. LAWR. Proc. Boston Soc. xiv. 1872, 303. (Socorro I.)
- 655. Phaëthon wthereue, Linn.—Cf. Freke, Sci. Proc. Roy. Dubl. Soc. 1879.\* (Banks of Newfoundland, Angust, 1876.)
- 665. Larus affinis, Reinh.—Vid. Medd. Nat. För. Kjobenhavn, 1853, 78. (Greenland.) 666a. Larus argentatus smithsonianus, Coues.—L. smithsonianus, Coues, Proc.
- Philad. Acad. 1862, 296. (North America.)
- 667. Larus caehinnans, Pall.—"Larus borealis, Brandt", BAIRD, Trans. Chicago Acad. i. 1869, 305. (St. Michael's, Alaska; Bischoff.)
- 671. Larus canus, Linn.—Cf. Brewer, Bull. Nutt. Orn. Club, iii. Jan. 1878, 50. (Labrador; Coues.)
- 689. Sterna Aleutica, Baird.—Trans. Chicago Acad. i. 1869, 321, pl. 31, fig. 1. (Kadiak, Alaska; Bischoff.)
- 692. Sterna anasthetu, Scopoli.—Sterna (Haliplana) anosthata, Coues, Key, 1872, 322 (Florida.)
- 694. Hydrochelidon leucoptera (Meisn.) Boie.—Cf. Brewer, Am. Nat. March, 1874, 188. (Lake Koshkonong, Wisconsin; T. Kumlien.)
- 705b. FULMARUS GLACIALIS RODGERSI (Cass.) Coues.—F. rodgersii, Cass. Proc. Philad. Acad, 1862, 290 (North Pacific); BAIRD, Trans. Chicago Acad. i. 1869, 323, pl. 34, fig. 1 (St. Georges I., Alaska).
- 710. Puffinus creatopus, Cooper.—Cf. Coues, Proc. Philad. Acad. 1864, 131. (Coast California.)
- 713. Puffinus guvia (Forst.) Finsch.—P. opisthomelas, Coues, Proc. Philad. Acad. 1864, 139. (Coast California.)
- Puffinus griseus (Gmel.) Finsch.—Nectris amaurosoma, Coues, Proc. Philad. Acad. 1864, 124.
- 716. Puffinus tenuirostris, Temm.—Neetris tenuirostris, Dall & Bannister, Trans. Chicago Acad. i. 1869, 303. (Kotzbue Sound.)
- Æstrelata bulweri (Jard.) Gigl. & Salvad.—Thalassidroma bulweri, Newton, Man. Nat. Hist. Greenl. 1875, 108.
- 720. HALOCYPTENA MICROSOMA, Coues.—Proc. Philad. Acad. 1864, 78. (Coast of California.
- 725. CYMOCHOREA HOMOCHROA, Coues.—Proc. Philad. Acad. 1864, 77. (Coast of California.)
- 737. Colymbus adamsi, Gray.—Proc. Zoöl. Soc. Lond. 1859, 167. (Alaska.)
- 763a. Lomvia troile Californica (Bryant) Cones.—Catarractes californicas, Bryant, Proc. Boston Soc. N. H. (Farallone Islands.)
- 764. Lomvia arra, Pall.—Cephus arra, Pall. Zoög. Rosso-As. ii. 1811, 347. (Alaska.)
- c. List of North American genera which have been described or added to the fauna since 1859, together with those whose names or orthography have been changed since that date.
  - HYLOCICHLA, Baird, Review Am. B. i. June 2, 1864, 12 (type, Turdus mustelinus, Gmel.).
    - The *Turdus musicus* of Europe is a strict congener of *T. mustelinus*, and it is possible that some generic name may have been based upon it previous to the imposition of *Hylocichla*.
  - 6. Turdus, Linnæus, S. N. ed. 10. i. 1758, 168 (type, T. viscivorus, Linn.).
    - The *T. iliacus*, although not agreeing strictly with *T. viscivorus* in details of external structure, is still hardly different enough to entitle it to separation. It is much more like true *Turdus* than either of the other American genera (*Hylocichla* and *Merula*).

<sup>\*</sup>We cannot cite the page of the "Proceedings", but the record appears on p. 44 of separate pamphlet entitled "A Comparative List of Birds found in Europe and North America. By Percy Evans Freke."

7. Merula, "Leach, 1816" (type, Turdus merula, Linn.).

The proper generic division of the typical thrushes is a matter of considerable difficulty. Of the North American generic groups, Hylocichla and Hesperocichla of Baird are sufficiently isolated, the latter being represented by a single species only, the former by all the smaller spotted species, besides the Song Thrush (Turdus musicus, Linn.) of Europe. I find no American species agreeing at all closely with Turdus viscivorus (the type of Turdus) in form; and a generic division based whelly or chiefly on coloration being out of the question, I find no other alternative than to adopt for the Robin and other American thrushes usually referred to "Planesticus" of Bonaparte (1854) the name Merula, Leach (1816), there being no essential difference in form between the type of the latter, Turdus merula, Linn. (Merula nigra, Leach) and our Robin (T. migratorius, Linn.); while a number of the Neotropical species exhibit the same sexual difference in coloration as T. merula. I would also refer to Merula the following Old World forms: Turdus pilaris, Linn. (type of Areeuthornis, Kaup, 1829), Turdus atrogularis, Temm. (type of Cichloides, Kaup, 1829), Turdus torquatus, Linn. (type of Thoracocincla, Reich., 1850), with perhaps some others.

- HESPEROCICHLA, Baird, Review Am. B. i. July, 1864, 32 (type, Turdus navius, Gmel.).
- 12. Galeoscoptes, Cabanis, Mus. Hein. i. 1850, 82 (type, Muscicapa carolineusis, Linn.).
- Cinclus, Bechstein, Gemein. Naturg. 1802 (type, Sturnus cinclus, Linn.). [Cf. BAIRD, Review Am. B. i. 1864, 59, foot-note.]
- 20. Cyanecula, Brehm, Vög. Deutschl. 1828 (type, Motacilla suecica, Linn.).
- 34. Phylloscopus, Boie, Isis, 1826, 792 (type?)
- Auriparus, Baird, Review Am. B. i. Aug. 1864, 85 (type, *Ægithalus flaviceps*, Snndev.).
- 61. Thryomanes, Sciater, Cat. Am. B. 1861, 22 (type, Troglodytes bewicki, Aud.).
- 65. Auorthura, Rennie, Montagn's Orn. Diet. 2d ed. 1831, 570 (type, A. communis, Rennie = Motacilla troglodytes, Linn.).
- 67. Telmatodytes, Cabanis, Mus. Hein. i. 1850, 78 (type, Certhia palustris, Wils.).
- 69. Motacilla, Linnieus, S. N. 1735 (type, M. alba, Linn.).
- 70. Budytes, Cuvier, Règ. An. i. 1817, 371 (type, Motacilla flava, L.).
- Helonwa.—Helinaia, Andubon, Synop. 1839, 66 (type, Sylvia swainsoni, Aud.).
   [Orthography emended by Agassiz, Nomencl. 1847. Cf. Newton, P. Z. S. 1879, 559. 7
- 90. Perissoglossa, Baird, Review Am. B. i. 1864, 180 (type, Motacilla tigrina, Gm.).
- 92. Peucedramus, Coues, in Henshaw's Orn. Wheeler's Survey, 1875, 201 (type, Sylvia olivacea, Giraud).
- 115. Siurus.—Cf. Coues, Bull. Nutt. Orn. Club.
- 124. Wilsonia, Bonaparte, Comp. List. 1838, 23 (type, Sylvia mitrata, Aud.?). [Cf. Coues, Bull. Nutt. Orn. Club, April, 1880, 95.\*]
- 131. Cardellina, "Dubus", Bonap. Consp. i. 1850, 312 (type, Cardellina amieta, Dubus = Muscicapa rubrifrons, Giraud).
- Ergaticus, Baird, Review Am. B. i. May, 1865, 264 (type, Sclophaga rubra, Swains.).
- 133. Basileuterus, Cabanis, in Schomb. Guiana, iii. 1848, 666 (type, Sylvia vermivora, Vieill.).
- 135. Vireosylria, Bonaparte, Comp. List. 1838, 26 (type, Muscicapa olivacca, Linu.).
- 140. Lanivireo, Baird, Review Am. B. i. May 23, 1866, 345 (type, Virco flavifrons, Vieill.?).

<sup>\*</sup> It is exceedingly doubtful whether Wilsonia, Bp., should displace Myiodioctes, Aud. Bonaparte's name occurs first in a mere list, is used only as a heading for a subgeneric group, and is unaccompanied either by a diagnosis or an indication of type. Audubon, however, only a year later, in designating the same group of birds by the new generic term Myiodioctes, gave an excellent diagnosis of the generic characters. It appears to us that the slight difference of date in favor of Bonaparte's name is greatly overbalanced by the pains which Audubon took to duly characterize his genus, thus conforming to the requirements of nomenclatural laws, which Bonaparte failed to do.

- 148. Lanius, Linnæus, S. N. ed. 10, 1758, 93 (type, L. excubitor, Linn.). [Cf. Coues, Birds Colorado Val.i. 1878, 539.]
- 153. Petrochelidon, Cabanis, Mus. Hein. i. 1850, 47 (type, Hirundo melanogaster, Sw. = P. swainsoni, Scl.).
- 155. Tachycineta, Cabanis, Mus. Hein. i. 1850, 48 (type, Hirundo thalassina, Sw.).
- Cottle, Boie.—Cf. Wharton, The Ibis, Oct. 1879; Coues, Bull. Nutt. Orn. Club, April, 1880, 96.
- 158. Stelgidopteryx, Baird, B. N. Am. 1858, 312, in text (type, Hirundô serripennis, Aud.).
- 167. Pyrrhula, Brisson, Orn. 1760 (type, Loxia pyrrhula, Linn. = Emberiza coecinea, Sandb.).
- 172. Loxia, Linnieus, S. N. ed. 10, i. 1758, 171 (type, L. currirostra, Linn.).
- 181. Astragalinus, Cabanis, Mus. Hein. i. 1851, 159 (type, Fringilla tristis, Linn.).
- 187. Centrophanes, Kaup, Ent. Gesch. Eur. Thierw. 1829 (type, Emberiza lapponica, Linn.).
- 190. Rhynchophanes, Baird, B. N. Am. 1858, 432, in text (type, Pleetrophanes maccowni, Lawr.).
- 224. Amphispiza, Coues, Birds N. W. 1874, 234 (type, Emberiza bilineata, Cass.).
- 244. Zamelodia, Coues, Bull. Nutt. Orn. Club, v. April, 1880, 98 (type, Loxia ludoviciana, Linn.).
- 247. Passerina, Vieillot, Analyse, 1816, 30 (type, Tanagra eyanea, Linn.). [Cf. Coues, l. c. 96.]
- 253. Phonipara, Bonaparte, Consp. i. 1850, 494 (type, Loxia eanora, Gmel.).
- 254. Spiza, Bonaparte, Obs. Wils. Orn. 1825 (part); Spece. Comp. 1827, 47 (type, Emberiza americana, Gmel.!). [See anteâ, 3.]
- 279. Sturnus, Linnens, S. N. ed. 10, i. 1758, 167 (type, S. vulgaris, Linn.).
- 285. Gymnocitta.—Cf. Coues, Bull. Nutt. Orn. Club, April, 1880, 98.
- 289. Cyanocitta, Strickland, Ann. Mag. N. H. xv. 1845, 260 (type, Corvus cristatus, Linn.). [Cf. Coues, Bull. Nutt. Orn. Club, April, 1880, 98.]
- 291. Aphelocoma, Cabanis, Mus. Hein. i. 1851, 221 (type, Garrulus californicus, Vig.).
- 299. Alauda, Linnæus, S. N. ed. 10, i. 1758, 165 (type, A. arvensis, Linn.).
- 308. Pitangus, Swainson, Zool. Jour. iii. 1827, 165 (type?).
- 309. Myiozetetes, Selater, ex. Schiff.—Myiozeta, "Schiff.", Bp. Compt. Rend. xxxviii. 1854, (type?).—Myiozetetes, Scl. P. Z. S. 1859, 46.
- 310. Myiodinastes, Bonaparte, Compt. Rend. xxxviii. 1854, 657 (type ?).
- 331. Ornithion, Hartlaub, Jour. für Orn. 1853, 35 (type, O. incrme, Hartl.).
- 332. Pachyrhamphus, "Gray, 1838" (type, Pachyrhyuchus euvieri, Spix = Tityra viridis, Vieill.).
- 333. Hadrostomus, Cabanis, Mus. Hein. ii. Oct. 24, 1859, 84 (type, *Tityra atricapilla*, Vieill.).
- 334. Eugenes, Gould, Mon. Troch. pt. xii. 1856 (type, Trochilus fulgens, Swains.).
- 337. Calypte, Gould, Introd. Troch. oct. ed. 1861, 87 (type, Ornismya costa, Boure.).
- 342. Atthis, Reichenbach, Aufz. der Colib. 1853, 12 (type, Ornismya heloisæ, Less. & Delattr.).
- 343. Stellula, Gould, Introd. Troch. oct. ed. 1861, 90 (type, Trochilus calliope, Gould).
- 344. Calothorax, Gray, Gen. B. 1840, 13 (type, Cynanthus lucifer, Swains.).
- 345. Amazilia, Lesson.—Amazilis, Less. Ind. Gen. et Syn. du Gen. Troch. 1832, p. xxvii. (type, Orthorhynchus amazili, Less.).—Amazilia, Reich. Av. Syst. Nat. 1849, pl. 39.
- 347. Basilinna, Boie, Isis, 1831, 546 (type, Trochilus leucotis, Vieill.).
- 348. IACHE, Elliot, Synop. Troch. March, 1879, 234 (type, Cynanthus latirostris, Swains.).
- 349. Cypselus, Illiger, Prodr. 1811, 229 (type, C. apus, Linn.). [Cf. anteâ, 6.]
- 350. Cypseloides, Streubel, Isis, 1848, 366 (type, Cypselus fumigatus, Natt.). [Cf. Sclater, P. Z. S. 1865, 614.]
- 354. Caprimulgus, Linnæns, S. N. ed. 10, i. 1858, 193 (type, C. europæus, Linn.). [Cf. Proc. U. S. Nat. Mus. i. 1878, 143; ib. iii. 1880, 5.]

Proc. Nat. Mus. 80——15

- 355. Phalenopthus, Ridgway, Proc. U. S. Nat. Mus. iii. March 27, 1880, 5 (type, Caprimulgus nuttalli, Aud.).
- 366. Xenopicus, Baird, B. N. Am. 1858, 83, in text (type, Leuconerpes albolarvatus, Cass.).
- 394. Aluco, Fleming, Phil. Zool. ii. 1828, 236 (type, Strix flammea, Linn.). [Cf. Newton, Yarrell's Brit. B. ed. 4, i. 150; Ibis, v. 94-105.]
- 395. Asio, Brisson, Orn. i. 1766, 28 (type, Strix otus, Linn.).
- 397. Strix, Linnaus, S. N.ed. 10, i. 1758, 92 (type, S. stridula, Linn.). [Cf. Newton, Ibis, ser. iii. vi. 94-105.]
- 399. Ululu, Cuvier, Règ. An. i. 1817, 329 (type, Strix uraleusis, Pall.).

In History of North American Birds (vol. iii, pp. 28-30) I adopted for the Great Gray Owl (Strix cincrea, Gmel.) the subgeneric name Scotiaptex, Swaius., based upon this species, and on p. 8 of these Proceedings raised the same name to generic rank. A subsequent examination of Strix urdlensis, Pallas, however, reveals the fact that the latter is strictly congeneric with S. cincrea and S. lapponica, and having been made the type of a genns Ulula by Cuvier, in 1817, the latter name must take precedence over Scotiaptex, which was not founded antil 1831.

- 408. Spectyto, Gloger, Handb. Naturg. 1842, 226 (type, Strix cunicularia, Mol.).
- 411. MICRATHENE, Coues, Proc. Philad. Acad. 1866, 51 (type, Athene whitneyi, Cooper.).
- 412. Hierofalco, Cuvier, Règ. An. i. 1817, 312 (type, Falco candicans, Gm.).
- Æsalon, Kaup, Natürl. Syst. 1829, 40 (type, Falco æsalon, Gmel. = F. regulus, Pall.)
- 419. RHYNCHOFALCO, Ridgway, Proc. Boston Soc. 1873, 46 (type, Falco femoralis, Temm. = F fusco-carulesceus, Vieill.).
- 426. Elanoides, Vieillot, Nouv. Dict. xxiv. 1818, 101 (type, Falco furcatus, Linn.).
- 434. Antenor, Ridgway, Proc. Boston Soc. Nat. Hist. May, 1873, 63 (type, Falco unicinctus, Temm.).
- 444. Urubitinga, Lesson, Rev. Zool. 1839, 132 (no type!); Lafr. in d'Orb. Dict. Hist. Nat. ii. 1842, 786 (type, Falco urubitinga, Gm. = F. zonurus, Shaw).
- 446. ONYCHOTES, Ridgway, Proc. Philad. Acad. Dec. 1870, 142 (type, O. gruberi, Ridgw.).
- 450. Thrasuëtus, Gray, Proc. Zool. Soc. Lond. 1837, 108 (type, Falco harpyia, Linn.).
- 451. Haliwëtus, Savigny.—This is the original and correct orthography. [Cf. Grav,
  Handb. i. 1869, 16; Coues, Bull. Nutt. Orn. Club, Apr. 1880, —.]
- 453. PSEUDOGRYPHUS, Ridgway, in B. B. & R. Hist. N. Am. B. iii. Jan. 1874, 337, 338 (type, *Vultur californianus*, Shaw).
- 455. Catharista, Vieillot, Analyse, 1816, 21 (type, Vultur urubu, Vieill. = V. atvata, Bartr. Cf. Ridgw. Bull. Nutt. Orn. Club, April, 1880, 80).
- 463. Engyptila, Sundevall, Met. Nat. Av. Disp. Tent. 1872, 156 (type, Columba rufuxilla, Rich. & Bern.).
- 467. Geotrygon, Gosse, Birds Jam. 1847, 316, foot-note (type, G. sylvatica, Gosse = Columba cristata, Temm.).
- 469. Ortalis.—Cf. Wharton, The Ibis, Oct. 1879, 450. [= Ortalida, Merrem (false orthography).]
- 471. Canace, Reichenbach, Av. Syst. Nat. 1851 (type, Tetrao canadensis, Linn.).
- DICHROMANASSA, Ridgway, Bull. U. S. Geol. & Geog. Surv. Terr. iv. No. 1, Feb. 5, 1878, 246 (type, Ardea rufa, Bodd.).
- 492. Hydranassa, Baird, B. N. Am. 1858, 660, in text (type, Ardea ludoriciana, Gmel. = A. tricolor, Müll.).
- 499. Mycteria, Linnæus, S. N. i. 1758, 140 (type, M. americana, Linn.).
- 501. Eudocimus, Wagler, Isis, 1832, 1232 (type, Scolopax rubra, Linn.). [Cf. Elliot, Ibis, 1877, 482.]
- 503. Plegadis, Kaup, Skizz. Ent. Gesch. 1829, 82 (type, Tantalus falcinellus, Linn.). [Cf. Sclater & Salvin, Ibis, 1878, 112.]
- 505. Ajaja, Reichenbach, Handb. 1851, p. xvi. (type, Platalea ajaja, Linn. = P. rosea, Briss.). [Cf. RIDGWAY, Proc. U. S. Nat. Mus. iii. 1880, 10.]

- 512. Vanellus, Meyer, Vög. Deutschl. i. 1810, 10 (ex. Linn. 1735; type, Tringa vanellus Linn.).
- Oxycchus, Reichenbach, Av. Syst. 1853, Introd. p. xviii. (type, Charadrius vociferus, Linn.).
- 522. Ochthodromus, Reichenbach, l. c. (type, Charadrius wilsonius, Ord).
- 523. Podasocys, Coues, Proc. Philad. Acad. 1866, 96 (type, *Charadrius montanus*, Towns.).
- 524. Scolopax, Linnaus, S. N. ed. 10, i. 1758, 145 (type, S. rusticula, Linn.).
- 530. Arquatella, Baird, B. N. Am. 1858, 717 (type, Tringa maritima, Briinn.).
- 533. Actodromas, Kaup, Sk. Ent. Eur. Thierw. 1829, 37 (type, Tringa minuta, Leisl.).
- 539. Pelidua, Cuvier, Règ. An. 1817, 490 (type, Tringa alpina, Linn.).
- 547. Totanus, Bechstein, Nat. Deutschl. 1803 (type, Scolopax calidris, Linn.).
- 554. Machetes, Cuvier, Règ. An. 1817 (type, Tringa pugnax, Linn.). [Cf. Coues, Bull. Nutt. Orn. Club, Apr. 1880, 100.]
- 555. Bartrania, Lesson, Traité Ois. 1831, 553 (type, B. laticauda, Less. = Tringa longicauda, Bechst.). [Cf. Cones, I. c.]
- 564. Lobipes, Cuvier, Règ. An. 1817 (type, Tringa hyperborea, Linn.).
- 565. Steganopus, Vieillot, Enc. Meth. 1823 (type, S. tricolor, Vieill. = Phalaropus wilsoni, Sab.).
- 568. Parra, Liunæus, S. N. i. ed. 12, 1766, 259 (type, P. jacana, Linn.). [For generic characters and illustrations, see these Proceedings, vol. i. pp. 163, 167, pl. iii.]
- 578. Ionoruis, Reichenbach, Av. Syst. 1853, 21 (type, Fulica martiuica, Linn.).
- 586. Olor, Wagler, Isis, 1832, 1234 (type, Cygnus musicus, Bechst. = Anas cygnus, Linn.).
- 590. Cheu, Boie, Isis, 1822 (type, Anas hyperborea, Pall.).
- 598. Philacte, Bannister, Proc. Philad. Acad. Nov. 1870, 131 (type, Anas canagicus, Sewast.).
- 619. Clangula, Fleming, Philos. Jour. 1828 (type, Anas clangula, Linn.). [Cf. Dresser, B. Eur. pt. xlvi. Dec. 1875; Coues, Bull. Nutt. Orn. Club, April, 1880, 101.]
- 635. NOMONYX, Ridgway, Proc. U. S. Nat. Mus. iii. March 27, 1880, 15 (type, Anas dominica, Linn.).
- 642. Phalaerocorax, Brisson, Orn. 1760 (type, Pelecanus carbo, Linn.). [Cf. Sharpe, Cat. B. Brit. Mus. iii. 1877, 146, foot-note.]
- 696. Megalestris, Bonaparte, Consp. ii. 1856, 206 (type, Lavus catarractes, Linn. = Catharacta skua, Briinn.). [Cf. Coues, B. N. W. 1874, 603, 604, where, however, Buphagus, Mæhring, is adopted; but Mæhring's names being inadmissible,\* Megalestris, Bp., "strictly its only synonym" seems the only one available.
- Phæbetria, Reichenbach, Av. Syst. Nat. 1853. pl. 26, fig. 348 (type, Diomedea fuligiuosa, Gmel.).
- 704. Ossifraga, Hombron & Jacquinot, Compt. Rend. xviii. 1844, 356 (type, Procellaria gigantea, Gmel.).
- 706. Priocella, Hombron & Jacquinot, Compt. Rend. xviii. 1844, 357 (type, Procellaria glacialoides, Smith = P. tenuirostris, Aud.).
- 707. Priofinus, Hombron & Jacquinot, Compt. Rend. xviii. 1844, 355 (type, Procellaria cinerea, Gmel. (?) = P. melauura, Bonn.).
- 717. Estrelata, Bonaparte, Consp. ii. 1855, 188 (type, Procellaria hasitata, Temm.).
- 720. HALOCYPTENA, Coues, Proc. Philad. Acad. 1864, 78 (type, H. microsoma, Coues).
- 721. Procellaria, Linniens, S. N. ed. 12, i. 1766, 212 (type, P. pelagica, Linn.).
- 722. Oceanites, Keyserling & Blasius, Wirb. Eur. 1840, 238 (type, Procellaria oceanica, Kuhl.).

<sup>\*</sup>Cf. Coues, Bull. Nutt. Orn. Club, April, 1880, p. 100, sp. 437, Machetes pugnax.

- Cymochorea, Coues, Proc. Philad. Acad. 1864, 75 (type, Procellaria leucorrhoa, Vieill.).
- 726. Occanodroma, Reichenbach, Av. Syst. 1853, xviii. (type, Procellaria furcata, Gmel.).
- Zehmophorus, Cones, Proc. Philad. Acad. 1862, 229 (type, Podiceps occidentalis, Lawr.).
- 732. Dytes, Kaup, Syst. Ent. Eur. Thierw. 1829 (type, Colymbus cornutus, Gmel.).
- 734. Tachybaptes, Reichenbach, Syst. Nat. Av. 1853, pl. 2 (type, Colymbus minor, Gmel.). 742. Utamania, Leach, Syst. Cat. 1816 (type, Alca torda, Linn.). [Cf. Coues, Proc.
- 742. Utamania, Leach, Syst. Cat. 1816 (type, Alca torda, Linn.). [Cf. Coues, Proc Philad. Acad. 1868, 18, 19.]
- 743. Fraterenla, Brisson, Orn. 1760 (type, Alea arctica, Linn.). [Cf. Coues, Proc. Philad. Acad. 1868, 21.]
- 745. Lunda, Pallas, Zorgr. Rosso-As. 1811, 363 (type, Alca cirrhata, Pall.). [Cf. COUES, Proc. Philad. Acad. 1868, 26.]
- 746. Ceratorhina.—Cf. Coues, Key, 1872, 341. [= Cerorhinea, Bonap. Ann. Lyc. N. Y. ii. 1828, 428 (false orthography).]
- 748. Simorhynchus, "Merrem, —, 1819 (type, Alca cristatella, Pall. fide G. R. Gray)". [Cf. Coues, Proc. Philad. Acad. 1868, 35.]
- 750. Ciceronia, Reichenbach, Av. Syst. Nat. 1853, (type, Phaleris microceros, Brandt = Uria pusilla, Pall.).
- 752. Alle, Link, Beschr. Natur.-Samml. Univ. Rostock, 1806, 17 (type, A. nigricans, Link = Alca alle, Linn.). [Cf. Coues, Bull. Nutt. Orn. Club, iv. Oct. 1879, 244.]
- 753. Synthliborhamphus, Brandt, Bull. Acad. St. Petersb. ii. 1837 (type, Alca antiqua, Gmel.).
- 763. Lomvia, Brandt, Bull. Acad. St. Petersb. ii. 1837, 345 (type, Colymbus troille, Linn.). [Cf. Coues, Proc. Philad. Acad. 1868, 75.]
- d. Species included in the catalogue which have not yet (according to the records) actually been taken within the prescribed limits.

[The following species enumerated in the catalogue have not, to this date, been taken within the United States; but all are known to occur so near our southern border as to render it quite certain that their capture within our limits is but a question of time and investigation. There are also included in this category all the species which are peculiar to the islands of Socorro and Guadalupe and the peninsula of Lower California.]

- 8. MERULA CONFINIS. (Cape St. Lucas.)
- 14. Harporhynchus cinereus. (Cape St. Lucas.)
- 18. Harporhynchus Graysoni. (Socotto.)
- 31. Regulus obscurus. (Gnadalnpe.)
- 43. Parus meridionalis. (Highlands of Mexico.)
- 49. PSALTRIPARUS MELANOTIS.\* (Highlands of Mexico.)
- 55a. Certhia familiaris mexicana. (Highlands of Mexico.)
- 57. Campylorhynchus affinis. (Cape St. Lucas.)
- 58a. Salpinctes obsoletus guadalupensis. (Guadalupe.)
- 62. Thryomanes Brevicauda. (Guadalupe.)
- 89. PARULA PITIAYUMI INSULARIS. (Socorto.)
- 171. CARPODACUS AMPLUS. (Guadalupe.)
- 195. Passerculus guttatus. (Cape St. Lucas.)
- 223. Junco insularis. (Guadalupe.)
- 238c. Pipilo maculatus consobrinus. (Guadalupe.)
- 238d. Pipilo Maculatus Carmani. (Socotto.)
- 240a. Pipilo fuscus albigula. (Cape St. Lucas.)

<sup>\*</sup>Probably seen by me in August, 1868, in the East Humboldt Mts., Nevada. (Cf. Orn. 40th Parallel Exp. 1877, p. 415.)

267. ICTERUS WAGLERI. (Mexico.)

288. PSILORHINUS MORIO. (E. Mexico.)

291. APHELOCOMA ULTRAMARINA COUCHII. (E. Mexico.)

347. Basilinna xantusi. (Cape St. Lucas.)

363a. Picus scalaris lucasanus. (Cape St. Lucas.)

377a. Melanerpes formicivorus angustifrons. (Cape St. Lucas.)

380. Colaptes rufipileus. (Guadalupe.)

381. Momotus ceruleiceps. (E. Mexico.)

384. Trogon ambiguus. (Mexico.)

393. Conurus holochlorus brevipes. (Socotto.)

415. FALCO ALBIGULARIS. (Whole of tropical America.)

424. Polyborus lutosus. (Guadalupe.)

.653. Sula piscator. (Socorro.)

655. PILETHON ÆTHEREUS. (Newfoundland banks; Socorro?)

460. ZENÆDURA GRAYSONI. (SOCGITO.)

Of the following species given in the catalogue no specimens are known to have been taken within the limits of the United States, with the exception of those described and figured in Giraud's "Sixteen Species of Texan Birds" (1841).\*

- 59. Catherpes mexicanus (=Certhia albifrons, Girand).
- 130. Setophaga miniata (=Muscicapa derhami, Giraud).
- 132. Ergaticus Ruber (=Parus leucotis, Girand).
- 133. Basileuterus culicivorus (=Muscicapa brasieri, Giraud).
- 134. Basileuterus Bellii (=Muscicapa bellii, Giraud).
- 160. Euphonia elegantissima (=Pipra galericulata, Girand).
- 182b. Astragalinus psaltria mexicanus (=Fringilla texensis, Girand).
- 309. Myiozetetes texensis (=Muscicapa terensis, Girand).
- 314. Myiarchus Lawrencei (=Tyraunula lawreneii, Girand).
- 329. Empidonax fulvifrons (=Muscicapa fulvifrons, Giraud).
- c. Species (chiefly Palwarctic) which occur only as stragglers or visitants in Eastern North America, or which occur regularly only in Greenland and adjacent portions of the continent.
  - [6.] TURDUS ILIACUS. Accidental in Greenland.
- [69.] MOTACILLA ALBA. Accidental in Greenland.
- [178.] ÆGIOTHUS CANESCENS. Resident in Greenland.
- [279.] STURNUS VULGARIS. Accidental in Greenland.
- [412a.] HIEROFALCO GYRFALCO ISLANDUS. Resident in South Greenland.
- [416.] ÆSALON REGULUS. Accidental in Greenland.
- [422.] TINNUNCULUS ALAUDARIUS, Accidental in Greenland.
- [435.] BUTEO VULGARIS. Accidental in Michigan? [Cf. Maynard, Bull. Nutt. Orn. Club, i. 18.]
- 452. Hallæëtus albicilla. Resident in Greenland.
- [488.] ARDEA CINEREA. Accidental in Greenland.
- [506.] Hæmatopus ostralegus. Accidental in Greenland.
- [514.] Charadrius Pluvialis. Accidental in Greenland.
- 518. ÆGIALITIS HIATICULA. Breeding in Greenland and west of Cumberland Gulf.
- [524.] SCOLOPAX RUSTICULA. Accidental in Newfoundland and Eastern United States.
- [526.] Gallinago media. Casual in Greenland and Bermudas.

<sup>\*</sup> These specimens are now in the collection of the United States National Museum.

<sup>†</sup>Strictly pelagic birds, which are more or less numerous off the coast, are excluded from this and the following lists.

- [539.] Pelidna Alpina. Breeds in Greenland and Hudson's Bay Territory.
- [540.] Pelidna subarquata. Casual in Eastern North America (several records).
- [546.] Limosa Egocephala. Accidental in Greenland.
- [547.] Totanus glottis. Accidental in Florida.
- [551.] Rhyacophilus ochropus. Accidental in Nova Scotia.
- [554.] Machetes Pugnax. Casual in Eastern North America (several records).
- [561.] NUMENIUS PHÆOPUS. Accidental in Greenland.
- [573.] Porzana Maruetta. Accidental in Greenland.
- [577.] CREX PRATENSIS. Casual in Eastern North America, including Greenland.
- [586.] OLOR CYGNUS. Accidental in Greenland.
- [587.] OLOR MINOR. Casual (?) in fur countries.
- [593.] ANSER ALBIFRONS. Breeds in South Greenland.
- [597.] Bernicla Leucopsis. Casual in Eastern North America.
- [611.] NETTION CRECCA. Casual in Eastern North America.
- 627. SOMATERIA MOLLISSIMA. Resident in Greenland and west side of Cumberland Gulf.
- [631.] MELANETTA FUSCA. Accidental in Greenland.
- [665.] Larus affinis. Accidental in Greenland.
- [694.] Hydrochelidon Leucoptera. Accidental in Wisconsin.
- [711.] PUFFINUS ANGLORUM. Casual (?) off Atlantic coast.
- [717.] ŒSTRELATA HÆSITATA. Accidental off Atlantic coast of U. S. .
- [718.] ŒSTRELATA BULWERI. Accidental near Greenland.
- 728. Fregetta Grallaria. Accidental off coast of Florida.
- [733.] Dytes auritus. Breeds in South Greenland.
- 743a. Fratercula arctica glacialis. Resident in Greenland.

## f. Palwarctic and oceanic species occurring only in Alaska and other parts of the Pacific coast.

- [20.] Cyanecula suecica. St. Michael's, Alaska, June 5, 1850. (See p. 215.)
- [34.] Phylloscopus Borealis. Breeds in Alaska.
- [44.] PARUS CINCTUS. Abundant resident in Alaska.
- [70.] BUDYTES FLAVA. Breeds abundantly in Alaska.
- [167..] Pyrrhula Cassini. Resident (?) in Alaska.
- [399a.] Ulula cinerea Lapponica. Casual (?) in Alaska (St. Michael's).
- [407a.] Surnia funerea ulula. Casual (?) in Alaska (St. Michael's).
- 511. APHRIZA VIRGATA. Casual along entire Pacific coast of America.
- [515a.] Charadrius dominicus fulvus. Regular autumnal visitant to Alaska.
- [519.] ÆGIALITIS CURONICA. Accidental in California (?).
- [533.] ACTODROMAS ACUMINATA. Accidental on coast of Alaska.
- 544. LIMOSA LAPPONICA NOVÆ-ZEALANDIÆ. Abundant visitant to Alaska.
- 553. Heteroscelus incanus. Whole Pacific coast. (Breeds.)
- [562.] NUMENIUS TAHITIENSIS. Accidental in Alaska (Kadiak).
- [702.] DIOMEDEA CULMINATA. Accidental off mouth of Columbia River (Audubon).
- [703.] Phœbetria fuliginosa. Casual off Pacific coast.
- [704.] Ossifraga gigantea. Accidental off Pacific coast of U.S.
- 706. PRIOCELLA TENUIROSTRIS. Casual (?) off Pacific coast.
- 707. PRIOFINUS MELANURUS. Accidental off coast of California.
- 710. Puffinus creatopus. Accidental (?) off coast of California.
- 713. Puffinus Gavia. Casual (?) off coast of Lower California.
- 715. Puffinus Griseus. Casual (?) off coast of Lower California.
- 716. Puffinus tenuirostris. North Pacific (casual?).
- [719.] DAPTION CAPENSIS. Accidental off coast of California.

g. Palwarctic species occurring both in Greenland and Alaska, but not recorded from any intermediate point in North America.

Catalogue No.

- [21.] SANICOLA ŒNANTHE. Breeds in Greenland and on west side of Cumberland Gulf.
- [72.] Anthus Pratensis. Accidental (?) in Greenland and Alaska.
- [512.] VANELLUS CRISTATUS. Accidental in Greenland (and Alaska?).
- [606.] Mareca Penelope. Occurs in various parts of North America, south to North Carolina, and San Francisco.
  - h. Tropical American species occurring only in southern portions of United States.

## EASTERN PROVINCE, INCLUDING FLORIDA AND COAST OF TEXAS.\*

- 137. VIREOSYLVIA CALIDRIS BARBATULA. (South Florida.) Hab. Cuba.
- 159. Certhiola Bahamensis. (Indian Key.) Hab. Bahamas.
- 184. Chrysomitris notata. (Accidental in Kentucky, fide Audubon.) Hab. high-lands of Mexico and Guatemala.
- 253. Phonipara Zena. (Key West.) Hab. West Indies in general.
- 265. ICTERUS VULGARIS. (South Carolina, etc.) Hab. Jamaica and northern South America.
- 302. Milvulus tyrannus. (Accidental in Mississippi, New Jersey, etc.) Hab. whole of tropical South America east of the Andes, Atlantic coast region of Central America (and Mexico?).
- 303. Tyrannus dominicensis. (Florida.) Hab. whole of West Indies.
- 357b. Chordelles popetue minor. (Florida.) Hab. Cuba and Jamaica.
- 386. Coccyzus seniculus. (Florida, Louisiana?) Hab. West Indies and parts of northern South America.
- 389. Crotophaga ani. (Tortugas; near Philadelphia!) Hab. West Indies and parts of northern South America.
- 420a, Tinnunculus sparverius isabellinus.
- 421. Tinnunculus sparverioides. (Florida.) Hab. Cuba.
- 429. Rostrhamus sociabilis plumbeus. (Florida.) Hab. Tropical America in general.
- 458. COLUMBA LEUCOCEPHALA. (Florida Keys.) Hab. West Indies; Honduras.
- 462. Zenæda amabilis. (Florida Keys.) Hab. Greater Antilles.
- 467. Geotrygon Martinica. (Florida Keys.) Hab. West Indies.
- 468. STARNŒNAS CYANOCEPHALA. (Florida Keys.) Hab. Cuba.
- 502. EUDOCIMUS RUBER. (Louisiana?) Hab. Northern South America; Jamaica.
- 503. PLEGADIS FALCINELLUS. (Florida, straggling northward.) Hab. Eastern Hemisphere chiefly.
- 578. IONORNIS MARTINICA. (Southern portions in general, straggling northward.)

  Hab. whole of tropical America.
- 581. Aramus Pictus. (Florida.) *Hab.* West Indies and Atlantic coast of Central America.
- 585. PHENICOPTERUS RUBER. (Florida Keys.) Hab. West Indies and shores of Gulf of Mexico and Caribbean Sea; Galapagos.
- 635. NOMONYX DOMINICUS. (Accidental on Lake Champlain and in Wisconsin.)

  Hab. whole of tropical America.
- 692. Sterna anæstheta. (Florida.) Hab. tropics generally.
- 734. TACHYBAPTES DOMINICUS. (Lower Rio Grande, in Texas.) Hab. Tropical America in general.

#### SOUTHWESTERN BORDER-TEXAS TO CALIFORNIA.

[The species of this list which are peculiar to the more elevated portions of Mexico and Gnatemala (including the contiguous southern border of the United States) are distinguished by an asterisk (\*) prefixed to the number. The avi-fauna of temperate Mexico is decidedly more nearly related to that of the Western Province of North America than to the tropical fauna of the Mexican tierra caliente or hot coast-region. The genera of Neotropical affinities are printed in italics.]

- \* 13a. Harporhynchus rufus longirostris. (Lower Rio Grande.)
- \* 15. Harporhynchus curvirostris. (Lower Rio Grande.)
- \* 26. Phainopepla nitens. (Texas to California.)
- \* 37. LOPHOPHANES ATROCRISTATUS. (Lower Rio Grande.)
- \* 39. LOPHOPHANES WOLLWEBERI. (New Mexico; Arizona.)
- \* 49. Psaltriparus melanotis. (Nevada?)
- \* 50. Auriparus flaviceps. (Texas to Arizona.)
- \* 56. Campylorhynchus brunncicapillus. (Texas to California.)
- \* 59. Catherpes mexicanus. (Lower Rio Grande?)
- \* 60a, Thryothorus Ludovicianus Berlandieri. (Lower Rio Grande.)
- \* 61b. Thryomanes bewicki leucogaster. (Lower Rio Grande to Arizona.)
- \* 83. Helminthophaga lucle. (Arizona.)
- \* 89a. Parula insularis nigrilora. (Lower Rio Grande.)
- \* 92. Peucedramus olivaceus. (Lower Rio Grande? Arizona.)
- \*104. Dendræca graciæ. (Arizona.)
- \*106. Dendreca chrysoparia. (Texas.)
- \*129. Setophaga picta. (Lower Rio Grande? Arizona.)
- \*130. Setophaga miniata. (Lower Rio Grande?)
- \*131. CARDELLINA RUBRIFRONS. (Lower Rio Grande? Arizona.)
- \*132. Ergaticus Ruber. (Lower Rio Grande?)
- 133. Basileuterus culicirorus. (Lower Rio Grande?)
- \*134. Basileuterus bellii. (Lower Rio Grande?)
- \*136. Vireosylvia agilis flavo-viridis. (Lower Rio Grande?)
- \*142. VIREO ATRICAPILLUS. (Texas.)
- \*147. VIREO VICINIOR. (Arizona; Southern California.)
- 160. Euphonia elegantissima. (Lower Rio Grande?)
- \*163. Pyranga hepatica. (New Mexico; Arizona.)
- \*164a. Pyranga Æsiva cooperi. (New Mexico; Arizona.)
- \*182a. Astragalinus psaltria arizonæ. (Upper Rio Grande to Arizona.)
- 182b. Astragalinus psaltria mexicanus. (Lower Rio Grande?)
- \*215. SPIZELLA ATROGULARIS. (Lower Rio Grande to Lower California.)
- \*222. Junco cinereus. (Arizona.)
- \*227. Peucæa arizon.e. (Lower Rio Grande to Arizona.)
- \*230a. Peucæa ruficeps boucardi. (Arizona.)
- \*236. Embernagra rufivirgata. (Lower Rio Grande.)
- \*241. PIPILO ABERTI. (Arizona.)
- \*242a. Cardinalis virginianus igneus. (Arizona; Lower California.)
- \*243. Pyrrhuloxia sinuata. (Lower Rio Grande to Lower California.)
- \*250. Passerina versicolor. (Lower Rio Grande.\*)
- 252. Spermophila moreletii. (Lower Rio Grande.)
- \*258a. Molothrus ater obscurus. (Texas to Lower California.)
- 259. MOLOTHRUS ÆNEUS. (Lower Rio Grande.)
- 263a. Sturnella Magna Mexicana. (Lower Rio Grande.)

- \*266. Icterus audubonii. (Lower Rio Grande.)
- \*268. ICTERUS PARISORUM. (Texas to Lower California.)
- \*269. ICTERUS CUCULLATUS. (Texas to Lower California.)
- \*275. Quiscalus macrurus. (Lower Rio Grande.)
- \*276. Quiscalus palustris. (Head of Gulf of California? coast of Louisiana.)
- \*295. APHELOCOMA SORDIDA ARIZONÆ. (Arizona.)
- \*296. Xanthura luxuosa. (Lower Rio Grande.)
- \*334. Eugenes fulgens. (Arizona.)
- \*342. Atthis heloise. (Texas.)
- \*344. Calothorax lucifer. (Arizona.)
- 345. Amazilia fuscicandata. (Lower Rio Grande.)
- \*346. Amazilia yucatanensis. (Lower Rio Grande.)
- \*348. IACHE LATIROSTRIS. (Arizona.)
- 356. NYCTIDROMUS ALBICOLLIS. (Lower Rio Grande.)
- \*358. Chordeiles acutipennis texensis. (Texas to Lower California.)
- \*363. Picus scalaris. (Texas to Arizona.)
- \*365. Picus stricklandi. (Arizona.)
- \*373. CENTURUS AURIFRONS. (Lower Rio Grande.)
- \*374. Centurus uropygialis. (Arizona.)
- 383. CERYLE AMERICANA CABINISH. (Texas to Arizona.)
- 390. CROTOPHAGA SULCIROSTRIS. (Lower Rio Grande.)
- \*391. Rhynchopsitta pachyrhyncha. (Rio Grande Valley?)
- \*402b. Scops asio maccalli. (Texas.)
- \*403. Scops trichopsis. (New Mexico; Stockton, Cal.?)
- \*404. Scops flammeolus. (North to about 40° in higher western mountains.)
- 410. GLAUCIDIUM PHALENOIDES. (Texas and Arizona.)
- \*411. MICRATHENE WHITNEYI. (Arizona; S. E. California.)
- 419. Rhynchofalco fusco-cærulescens. (Texas; New Mexico.)
- 434. Antenor unicinctus harrisi. (Louisiana to Lower California.)
- 440. Buteo abbreviatus. (Arizona; Southern California.)
- 441. Buteo albicaudatus. (S. Texas.)
- 444. Urbitinga anthracina. (Arizona.)
- 445. ASTURINA NITIDA PLAGIATA. (Arizona.\*)
- 450. Thras.etus harpyia. (Lower Rio Grande; Louisiana?)
- 457. COLUMBA ERYTHRINA. (S. Texas.)
- \*463. Engyptila albifrons. (S. Texas.)
- 464. MELOPELIA LEUCOPTERA. (Texas to Lower California.)
- \*466. Scardafella inca. (Southern Texas.)
- \*469. Ortalis vetula maccalli. (S. Texas.)
- \*470. MELEAGRIS GALLOPAVO. (New Mexico; Upper Rio Grande in Texas.)
- 483. LOPHORTYX GAMBELI. (W. Texas to Arizona.)
- 484. Callipepla squamata. (W. Texas to Arizona.)
- 485. CYRTONYX MASSENA. (W. Texas to Arizona.)
- 499. Mycteria americana. (Southern Texas.)

## ENTIRE SOUTHERN BORDER.

- 423. Polyborus Cheriway.
- 427. Elanus glaucus.
- 455. Catharista atrata.
- 465. CHAMÆPELIA PASSERINA.

<sup>\*</sup>Accidental in Southern Illinois (only once observed).

i. Supposed valid species described by Audubou and Wilson, which have not since been met with, and of which no specimens are known to exist in collections.

Catalogue No.

- REGULUS CUVIERI, AUD. Orn. Biog. i. 1832, 288, pl. 55 ("Banks of Schuylkill River, Pa., June, 1812").—Baird, B. N. Am. 1858, 228; Review, i. 1864, 66.—B. B. & R. Hist. N. Am. B. i. 1874, 75, pl. 5, fig. 7.
- Perissoglossa Carbonata (Aud.) Ridgw.—Sylvia carbonata, Aud. Orn. Biog. i. 1831, 308, pl. 60 (Kentucky).—Dendroica carbonata, Baird, B. N. Am. 1858, 287; Review, i. 1865, 207.—Perissoglossa carbonata, B. B. & R. Hist. N. Am. B. i. 1874, 214, pl. 12, fig. 3.
- 112. Dendræca Montana (Wils.) Baird.— 'ylvia montana, Wils. Am. Orn. v. 1812, 113, pl. xliv. fig. 2 ("Blue Mts. of Pennsylvania").—Aud. Orn. Biog. v. 294 ("California").—Dendroica montana, Baird, B. N. Am. 1858, 279; Review, i. 1865, 190.—Dendræca montana, B. B. & R. Hist. N. Am. B. i. 1874, 271.
- WILSONIA MINUTA (Wils.) Bp.—Muscicapa minuta, WILS. Am. Orn. vi. 1812, 62,
   pl. l. fig. 5.—Aud. Orn. Biog. v. pl. 434, fig. 3; B. Am. i. pl. 67.—Myiodioetes minutus, Baird, B. N. Am. 1858, 293; Review, 1865, 241.—B. B. & R. Hist. N. Am. B. i. 1874, 316, pl. 16, fig. 2.
  - 1. List of untenable species and races of North American birds described since 1858.
  - Helminthophaga ruficapilla var. ocularis, B. B. & R. Hist. N. Am. B. i. 1874, 191.
     (Chicago, Ill.) = No. 85.
  - (?) 2. Helminthophaga ruficapilla var. gutturalis, B. B. & R. Hist. N. Am. B. i. 1874, 191. (East Humboldt Mts., Nevada; Ft. Tejon, Cal.) = No. 85?
  - 3. Helminthophaga celata var. obscura, B. B. & R. Hist. N. Am. B. i. 1874, 192. (Georgia and Florida.) = No. 86.
  - 4. Hirundo bicolor var. respertina, Cooper, Am. Nat. x. Feb. 1876, 91. (California.) = No. 155.
  - Collyrio chemingensis, Gregg, Proc. Elmira Acad. i. 1870, 9. (New York.) = No. 148, juv.
  - Hesperiphona respertina var. montana, Ridgw. in B. B. & R. Hist. N. Am. B. i. 1874, 449, pl. 22, fig. 4. (Southern Rocky Mts. and mountain regions of Mexico.) = No. 165.
  - (?) 7. Loxia atrata, von Homeyer, Jour. für Orn. 1879, 179. (North America.) = No. 173?? \*
  - 8. Leucosticte campestris, BAIRD, Orn. Cal. i. 1870, 163. (Colorado.) = No. 175a (individual variation).
  - (?) 9. Linaria brunnescens, von HOMEYER, Jour. für Orn. 1879, 184. ("Lapland, Grönland, Schweden.")† = No. 179a?

"This is decidedly not a melanism, as is shown, not alone by the character of the coloration, but by the courrence of two old males exactly alike."

"My two birds came from North America."-[Translation.]

<sup>\* &</sup>quot;Somewhat smaller than L. leucoptera, but with about the same length of wing, the tail at least 10mm shorter. Distinguished by its coloration. The red in ground tone is nearest that of the "Hakengimpel" [Pinicola enucleator] but everywhere darker, and saturated, as it were, with black, this color extending indefinitely both on the back as well as on the middle of the belly to the breast, as also on the lower tail-coverts. The feet are even darker colored than in leucoptera. The wing-bands are unaffected for the most part, although not quite so broad. This is especially seen in the first or terminal band, which begins, scarcely visible, at the edge of the wing, and remains very inconsiderable to the middle, then quickly expands to a great rounded spot. The extent of the second band is everywhere much more limited than in leucoptera."

<sup>†&</sup>quot;The dark Linnet is nearest to this species in size, but it is easily distinguished by many features. The whole upper parts are blackish brown, with very narrow margins to the feathers of whitish and rusty yellow, which, on the rump, return to the ground color; on the sides of the head the feathers also have very fine, rusty white margins; the throat-spot is very large; the entire under parts to the middle of the belly are densely covered with many large, dark brown longitudinal streaks. Two birds killed in Lapland, June-July (sexes not determined), show a tender red on the side of the head. The

Catalogne No.

- Ægiothus fuscescens, Coues, Proc. Philad. Acad. 1861, 222. (Labrador.) = No. 179 (midsummer plumage).
- 11. Egiothus rostratus, Coues, l. c. (Greenland.) = No. 179a (midsummer plumage).
- 12. Centronyx ochrocephalus, Aiken, Am. Nat. vii. 1873, 237. (El Paso Co., Colorado.) = No. 191 (autumnal plumage).
- Passerenlus caboti, B. B & R. Hist. N. Am. B. iii. 1874, pl. xlvi. fig. 9. (Nahant, Massachusetts.) = No. 233. [Not described!]
- Spizella evura, Coues, The Ibis, 1865, 118, 164. (Ft. Whipple, Ariz.) = No. 215, young.
- Passerella obscura, Verrill, Proc. Boston Soc. ix. 1862, 153. (Anticosti I.)
   No. 235, young.
- 16. Hedymeles melanocephalus var. capitalis, B. B. & R. Hist. N. Am. B. ii. 1874, 74. (Pacific coast of Mexico and United States.) = No. 245.
- Guiraca carulea var. eurhyncha, Coues, Am. Nat. viii. 1874, 563. (Mexico.) = No. 246.
- Dolichonyx oryzivorus var. albinucha, Ridgw. Bull. Essex Inst. v. Nov. 1873, 192.
   (Missouri plains to Salt Lake Valley.) = No. 257.
- Empidonax pygmæus, MINOT, Land and Game Birds New England, 1877, —. (Near Boston, Mass.) [Aris fictita!]
- Dryobates hyloscopus, Cab. & Heine, Mus. Hein. iv. June 25, 1863, 69. (San José, Cal.) = No. 360b.
- 21. Dryobates homorus, Cab. & Hein. Mus. Hein. iv. June 25, 1863, 65. (California.) = No. 361a.
- Picus curieri, Malli. Mon. Pic. i. 1861, 85, pl. 22, fig. 3. (North America.) = No. 360, Q ud.
- 23. *Picns turati*, Malh. Mon. Pic. i. 1861, 125, pl. 29, figs. 5, 6, 7. (California and Rocky Mountains.) = No. 361, 9 ad.
- Chamapelia passerina var. pallescens, Baird, Proc. Philad. Acad. 1859, 305.
   (Cape St. Lucas.) = No. 465.
- 25. Pediocætes kennicotti, Suckl. Proc. Philad. Acad. 1861, 361. = No. 478.
- 26. Bonasa jobsii, Jaycox, Am. Nat.
- Ibis thalassinus, Ridgw. Am. Nat. viii. Feb. 1874, 110. (Pacific coast of America, from California to Chili.) = No. 504, jnv.
- 28. Ardea cyunirostris, Cory, Birds of the Bahama Islands, 1880, —. (Bahamas.) = No. 492, breeding plumage.
- Cygnus passmorci, Hincks, Pr. Linn. viii. 1864, 1. (Toronto, Canada.) = No. 589, juv.
- 30. Bernicla barnstoni, Ross, Canad. Nat. vii. April, 1862, —. = 594, var.?
- 31. Berniela lencolama, Murry, Edinb. Phil. Jour. April, 1859, 226, pl. 4, fig. 1. = 594, yar.
- 32. Pelecanus occipitalis, Rigdw. Am. Sportsman, iv. 1874, 297. (Nevada.) = No. 640, adult, breeding plumage, after loss of occipital crest, the latter replaced by dusky-grayish patch.
- Thalasseus cuspius var. imperator, Coues, Proc. Philad. Acad. 1862, 538, in text. (North America.) = No. 680.
- 34. Sterna portlandica, Ridgw. Am. Nat. viii. 1874, 433. (Portland, Maine.) = No. 687, juv., second year
- Sterna fuliginosa var. crissalis, Baird, Pr. Boston Soc. xiv. 1872, 285. (Socorro I., N. W. Mexico.) = No. 681.

wing-bands are merely indicated. The bill is very characteristic. It is somewhat weaker at the base than in  $L.\ hornemanni$ , but longer and remarkably darker in all seven specimens.

<sup>&</sup>quot;The bird figured by Dresser on the second plate (lower figure) belongs here, and is by no means the young of L. hornemanni, as supposed by Dresser."

<sup>[</sup>This description accords well in every particular with the dark summer stage of Egiothus linaria holbölli, described in 1861 by Dr. Coues as E. rostratus, the type of which came from Greenland.—R. R.]

- k. List of exotic species which have been attributed to North America by various authors, but apparently without sufficient evidence of their occurrence.\*
  - 1. Anthus cervinus, Pall.—Zander, Jour. für Orn. Extraheft i. 1853, 64. (Aleutian Islands.)
  - Geothlypis equinoctialis (Gmel.) Caban.—Sylvia delafieldii, Aud. Orn. Biog. v. 1839, 307 ("Oregon").—Trichas delafieldii, Aud. B. A. Am. ii. 1841, 81, pl. 103.— Geothlypis velatus, BAIRD, B. N. Am. 1859, 243; Cat. 1859, No. 171.
  - 3. Lanius lahtora, Sykes.—Lanius elegaus, Swains. Faun. Bor. Am. ii. 1831, 122 (fur countries).—Nutt. Man. ii. 1832, 566. [Not Collurio elegans, Baird.]
  - Progue leucogastra, Baird.—P. chalubea, Cass. Illustr. 1856, 246 (California, fide J. G. Bell).
  - Astragalinus yarrelli (Aud.) Caban.—Carduelis yarrelli, Aud. Synop. 1839, 117 ("California"); B. Am. iii. 1841, 136, pl. 184.—Chrysomitris yarrelli, BAIRD, B. Am. 1858, 421; Cat. 1859, No. 312.
  - Astragaliuus barbutus (Mol.) ——.—Carduelis stanleyi, Aud. Synop. 1839, 118 ("California"); B. Am. iii. 1841, 137, pl. 185.—Chrysomitris stanleyi, BAIRD, B. N. Am. 1858, 420; Cat. 1859, No. 311.
  - HYPOLIA ARCTOA (Pall.) Ridgw.—Leucosticte arctoa, Caban. Mus. Hein. i. 1851, 154 ("Russich-America").—Leucosticte arctous, Baird, B. N. Am. 1858, 430; Cat. 1859, No. 324.
  - 8. CARPODACUS ILEMORRHOUS (Licht.) Scl.—BAIRD, B. N. Am. 1858, 417, foot-note (North America?); Cat. 1859, No. 309.
  - 9. Loxia pityopsittacus, Bechst.—Cf. Nuttall, Man. Orn. Land Birds, ed. 1832, 537 ("high northern regions of America", fide TEMMINCK).
  - "Zonotrichia" pileata (Bodd.) —.—Fringilla mortonii, Aud. Orn. Biog. v. 312;
     B. Am. iii. 1841, 152, pl. 190 ("North California").
  - Cynchramus schwuiclus (Linn.) Boie.—Emberiza schwuiclus, NUTT. Man. Orn. Land Birds, ed. 1832, ii. 586 ("vicinity of Harrisburg in Pennsylvania", fide Audubon).
  - Melanocorypha calandra (Linn.) Boie.—Alanda calandra, Linn., Sw. & Rich. F.
     B. A. ii. 1831, 244 ("fur countries"; spec. presented by the Hudson's Bay
     Co. said to be in the British Museum).—Nutr. Man. ii. 1832, 580.
  - Trupialis militaris (Linn.) Bp.—Baird, B. N. Am. 1858, 533 ("California"); Cat. 1859, No. 405.
  - 14. ICTERUS MELANOCEPHALUS (Wagl.) Gray.—Cass. Illustr. 1856, 137, pl. 21 (Texas and New Mexico).—Baird, B. N. Am. 1858, 543 (not given as North American!); Cat. 1859, No. 410.
  - CALOCITTA COLLIEI (Vig.) Finsch.—"Pica bullockii, Wagl.", Aud. B. Am. iv. 1842,
     105, pl. 229 ("woody portions of North California").—Garrulus bullockii,
     Nutt. Man. i. 1832, 230 ("Columbia R.").
  - Tyrannus melancholicus, Vieill.—BAIRD, B. N. Am. 1858, 176 (not given as North American); Cat. 1859, No. 129.
  - Lampornis violicanda (Bodd.) Elliot.—" Trochilus mango, Linn.", Aud. Orn. Biog. ii. 480; B. Am. iv. 1842, 186, pl. 251 ("Florida Keys").—Lampornis mango, Baird, B. N. Am. 1858, 130; Cat. 1859, No. 100.
  - CAMPEPHILUS IMPERIALIS (Gould) Gray.—Pieus imperialis, Aud. Orn. Biog. v. 313; B. Am. iv. 1842. 212 ("Rocky Mountains and North California).—Cass. Illustr. 1856, 285, pl. 49.—Baird, B. N. Am. 1858, 82; Cat. 1859, No. 73.

<sup>\*</sup>This list does not include American species wrongly supposed by authors to be the same as European species, and so named, e.g. Circus "eyaneus" for C. hudsonius, Regulus "cristatus" for R. satrapa, Troglodytes "parvulus" or T. "europeus" for T. hyemalis, etc., etc., but only those which were through actual error (as it appears) wrongly attributed to North America. Species which are most likely to have occurred in North America are printed in small capitals; those whose occurrence would in any case be purely accidental are printed in italics.

- Hylotomus scapularis (Vig.) Ridgw.—"Pieus lineatus, Linn.", Aud. Offi. Biog. v. 315; B. Am. iv. 1842, 233 ("Columbia River").
- STRIX STRIDULA, Linn.—S. aluco, NUTT. Man. i. 1832, 135 (Newfoundland and Hudson's Bay).
- CARINE NOCTUA (Scop.) Kaup.—"Strix passerina, LINN.", AUD. Orn. Biog. v. 269.—"Surnia passerina, Linn.", AUD. B. Am. i. 1840, 116 ("Pictou, Nova Scotia").
- Spectyto cunicularia (Mol.) ——.—Athene cunicularia, Cass. in Baird's B. N. Am. 1858, 60 ("North America, west of Rocky Mountains").
- 23. Thalassoltus pelagicus (Pall.) Kaup.—Aquila pelagica, Pall. Zoögr Rossolas. i. 1811, 343 (Russian America, fide Steller).—Haliaëtus pelagicus, Cass. Illustr. 1856, 31, pl. 6; in Baird's B. N. Am. 1858; Baird, Cat. 1859, No. 40.
- 24. Sareorhamphus gryphus (Linn.) Dum.—Cathartes gryphus, Bonap. Am. Orn. iv. 1833, 318, pl. 22.—Nutt. Man. i. 1832, 35.
- 25. GYPARCHUS PAPA (Liun.) Glog.—Cathartes papa, NUTT. Man. i. 1832, 40 ("from the 30th degree of north latitude to the 32d in the southern hemisphere").
- Cathartes Burrovianus, Cass. in Baird's B. N. Am. 1858, 6 ("Lower California"); Baird, Cat. 1859, No. 4. [Cf. Ridgway, Bull. Nutt. Orn. Club, v. April, 1880, 83.]
- 27. LOPHORTYX ELEGANS (Less.) Nutt.—Ortyx elegans, Nutt. Man. ed. 1840, i. 792 ("Upper California", fide Lesson).
- Butorides Brunnescens (Gundl.) Baird.—Baird, B. N. Am. 1858, 677 (in text);
   Cat. N. Am. B. 1859, No. 494.
- Hæmatopus ater, Vieill.—Hæmatopus townsendii, Aud. Orn. Biog. v. 1839, 247, pl. 427; B. Am. v. 1842, 245, pl. 326.—Hæmatopus ater, Baird, B. N. Am. 1858, 700; Cat. 1859, No. 514.
- 30. "TRINGA" PLATYRHYNCHA, Temm.—NUTT. Man. ii. 1832, 114 (Arctic America, fide Temminck and Bonaparte).
- Actodromas minuta (Linn.) Kaup.—Tringa minuta, Sw. & Rich. F. B. A. ii. 1831, 385 (Nelson and Hayes Rivers; "seen abundantly in the autumn").— NUTT. Man. ii. 1834, 119.
- Actodromas temmincki (Leisl.) Ridgw.—Tringa temmincki, Nutt. Man. ii. 1832, 119 (Arctic America).
- 33. Totanus calidris (Linn.) Bechst.—Sw. & Rich. F. B. A. ii. 1831, 391 ("Hudson's Bay"; spec. in British Museum).—Nutt. Man. ii. 1834, 155.
- 34. Heliornis fulica (Bodd.) ——.—H. surinamensis, NUTT. Man. ii. 1832, 510 ("accidental visitor in the Middle States of the Union").
- 35. Anser segetum (Gmel.) Eonap.—Nutt. Man. ii. 1832, 348 (Canada and Hudson's Bay).
- 36. Cairina Moschata (Linn.) Caban.—Anas moschata, Nutt. Man. ii. 1832, 403 Lower Mississippi and Gulf coast of U. S.).
- 37. ŒDEMIA NIGRA (Linn.) Hen.—Fuligula nigra, NUTT. Man. ii. 1832, 423 ("coast of the United States").
- 38. Mergellus Albellus (Linn.) Selby.—Mergus albellus, Wils. Am. Orn. iii. pl. lxxi. fig. 4 (New England and New York; nun.erous).—Nutt. Man. ii. 1832, 467.—Aud. Orn. Biog. iv. 350; B. Am. vi. 1843, 408, pl. 414 ("Lake Barataria, not far from New Orleans").
- 39. Phalacrocorax graculus (Linn.) Leach.—Nutt. Man. ii. 1832, 484 ("South of Greenland"; United States in winter).
- Phalacrocorax pygmaus, Pall.—Nutt. Man. ii. 1832, 487 (Northern North America, fide Bonaparte).
- 41. Phalacrocorax africanus (Gmel.) Dumont.—Nutt. Man. ii. 1832, 488 ("United States", fide Audubon).
- 42. Larus fuscus, Linn.—Nutt. Man. ii. 1832, 302 (Greenland, Newfoundland, and Hudson's Bay).

- 43. Larus capistratus, Temm.—Nutt. Man. ii. 1832, 290 (Delaware R. and Chesapeake Bay).
- 44. LARUS MINUTUS, Pall.—Sw. & RICH. F. B. A. ii. 1831, 426 (given on Sabine's authority).—NUTT. Man. ii. 1832, 289.—Chroicocephalus minutus, LAWR. in Baird's B. N. Am. 1858, 853.—BAIRD, Cat. 1859, No. 671.
- 45. DIOMEDEA EXULANS, Linn.—NUTT. Man. ii. 1832, 340 ("accidentally to the coasts of the central part of the Union").—LAWR. in Baird's B. N. Am. 1858, 821.—BAIRD, Cat. 1859, No. 630.
- Podiceps Cristatus (Linn.) Lath.—Sw. & Rich. F. B. A. ii. 1831, 410 (throughout fur countries).—Nutt. Man. ii. 1832, 250.—Lawr. in Baird's B. N. Am. 1858, 893.—Baird, Cat. 1859, No. 703.
- 47. Tachybaptes minor (Linn.) Coues.—*Podiceps minor*, Nutt. Man. ii. 1832, 257 (Hudson's Bay).
- 1. Partial list of forcign birds which have been introduced to the United States, and those which have been captured after escape from confinement.

#### SPECIES INTRODUCED WITH A VIEW TO THEIR NATURALIZATION.\*

- 1. Passer domesticus (Linn.) Leach. European House Sparrow. The attempted naturalization of this bird has proved decidedly successful. The case is so notorious that further comment is unnecessary.
- 2. Passer Montanus (Linn.) Stephens. European Tree Sparrow. Has become naturalized in the vicinity of Saint Louis, Mo., but the history of its introduction is unknown. (See Merrill, Bull. Nutt. Orn. Club.)
- Alauda arvensis, Linn. Skylark. Partially naturalized in the vicinity of Cineinnati, on Long Island, and perhaps other localities.
- 5. COTURNIX COMMUNIS (Liun.) Bonn. European Quail. Introduced to various local ities in the Eastern United States, and partially naturalized.

#### SPECIES WHICH HAVE BEEN CAPTURED AFTER ESCAPE FROM CONFINEMENT.

- AMADINA RUBRO-NIGRA, —. Brunswick, Maine, March, 1879; Leslie A. Lee. (Allen, Bull, Nutt. Orn. Club, April, 1880, 119.) Hab. India.
- 2. CRITHAGRA BUTYRACEA, —. South Scituate, Mass., in midwinter. (Brewer, Proc. Bost. Soc. xx. 271.) Hab. South Africa.
- 3. LIGURINUS CHLORIS (Linn.) Koch. Lowville, Lewis Co., N. Y., March 19, 1878; R. B. Hongh. (Cf. Bull. Nutt. Orn. Club, Apr. 1880, 119.) Hab. Europe.
- CARDUELIS ELEGANS, Steph. Eastern Massachusetts, many captures. (ALLEN, Bull. Nutt. Orn. Club, Apr. 1880, 120.) Hab. Europe.
- Serinus Meridionalis, Brehm. Western Massachusetts, in winter. (Allen, l. c.) Hab. Europe.
- 6. Corvus frugilegus, Linn. Washington, D. C., August, 1879.

An example of this species was seen by me in August, 1879, in the grounds of the Agricultural Department in Washington. It was perched in a maple tree near one of the outbuildings, was very tame, and flew laboriously, as if very recently escaped from confinement. I am, as yet, ignorant of the history of this specimen, nor have I since seen it.

- 7. Conurus xanthogenius, Bp. Hab. St. Thomas, West Indies.
  - An example of this species, shot in a grove near Washington, by Dr. D. W. Prentiss, is in the National Museum collection. Of course it was an escaped cage-bird.
- 8. CHENALOPEX .EGYPTIACA (Linn.) Steph. Carnarsie, Long Island, Jan. 3, 1877. (AKHURST, Bull. Nutt. Orn. Club, ii. Apr. 1877, 52.) Hab. Southern Europe and Africa.

<sup>\*</sup> This list does not include domesticated birds.

<sup>†</sup>This list is, of course, very incomplete; it includes merely a few species, the records of whose capture I happen to have at hand. A more complete list would be desirable, but want of time forbids its compilation in the present connection.

Table of families of North American birds, showing number of genera and species of each according to the foregoing catalogue.\*\*

	Number of the cata- logue.	Number of gen- era.	Number of species.†
1. Turdidæ 2. Cinclidæ	1-18 19	8	18 + 8 = 26
3. [Luscinidæ]	20	1	1
4. Šaxicolide	21-24	2	4
5. PTILOGONATIDE	25-26	$\frac{2}{3}$	2
6. Sylviidæ 7. Chamwidæ	27-34 35	1	$\frac{8}{1} + 1 = 9$
8. Paride	36-50	4	15 + 3 = 18
9. Sittidæ	51-54	1	4 + 1 = 5
10. Certhiidæ 11. Troglodytidæ	55 56-68	$\frac{1}{9}$	$\begin{vmatrix} 1+1=2\\ 13+9=21 \end{vmatrix}$
11. Troglodytidæ	69-73	4	$\begin{vmatrix} 13 + 9 = 21 \\ 5 \end{vmatrix}$
13. MNIOTILTIDÆ	74-134	18	61 + 8 = 69
14. VIREONIDÆ	135-147	3	13 + 3 = 16
15. Laniidæ	148-149 150-151	1	$\frac{2}{2} + 2 = 4$
17. Hirundinidæ	152-158	6	$\tilde{7} + 1 = 8$
18. Cærebidæ	159	1	1
19. TANAGRIDÆ	160-164	2	5 + 1 = 6
20. Fringillidæ	165-256 257-278	36 8	91 + 35 = 126 22 + 5 = 27
22. [Sturnidæ]	279	1	1
23. Corvidæ	280-298	9	19 + 7 = 26
24. [Alaudidæ]	299-300	$\frac{2}{11}$	2 + 2 = 4
25. Tyrannidæ 26. Cotingidæ	301 <b>-</b> 331 332 <b>-</b> 333	$\frac{11}{2}$	$\begin{vmatrix} 31 + 2 = 33 \end{vmatrix}$
27. Trochilide	.334-348	10	15
28. Cypselidæ	349-352	3	4
29. Caprimulgidæ	353-358	5	6 + 2 = 8
30. Picidæ 31. Momotidæ	359-380 381	10 1	22 + 10 = 32
32. Alcedinidæ	382-383	1	2
33. Trogonidæ	384	1	1
34. Cuculidæ	385-390	3 2	6 3
35. Psittacide 36. Strigidæ	391–393 394–411	12	18 + 11 = 29
37. Falconidæ	412-452	23	41 + 12 = 53
38. Cathartidæ	453-455	3	3
39. Columbidæ	456–468 469	10 1	13
40. Cracidæ 41. Meleagridæ	470	1	1 + 1 = 2
42. Tetraonidæ	471-479	$\bar{6}$	9 + 7 = 16
43. Perdicidæ	480-485	5	6 + 3 = 9
44. Ardeidæ 45. Ciconiidæ	486-498 499-500	11 2	$\begin{array}{c} 13 \\ 2 \end{array}$
45. Ciconiidæ 46. Ibididæ	501-504	2	4
47. Plataleidæ	505	1	1
48. Hæmatopodidæ	506-508	1	3
49. Strepsilidae 50. Charadriidae	509-511 512-523	2 7	$\begin{vmatrix} 3 \\ 12 + 2 = 14 \end{vmatrix}$
50. Charadriidæ 51. Scolopacidæ	524-562	21	39 + 4 = 43
52. Phalaropodidæ	563-565	3	3
53. Recurvirostride	566-567	2	2
54. Parridæ 55. Rallidæ	568 569–580	$\frac{1}{6}$	$\begin{vmatrix} 1 \\ 12 + 2 = 14 \end{vmatrix}$
56. Aramidæ	581	1	1 7 2 - 11
57. Gruidæ		1	3
	27 (1 4		-1: Balmanatia

<sup>\*</sup>Families peculiar to America in small capitals; peculiar to North America in italics. Palæarctic families not represented by a peculiar species in America are placed in brackets.

<sup>†</sup>The figures following the sign + denote the number of recognized races not distinguished by a separate number in the catalogue; those in the last column the total of species and races.

### Table of families of North American birds—Continued.

		Number of the cata- logue.	Number of gen- era.	Number of species.
58. Phenicopteridæ		585	1	1
59. Anatidæ		586-638	30	53 + 6 = 59
			1	1
			1	2
62. Phalacrocoracid	æ	642-648	1	7 + 3 = 1
63. Plotidæ		649	1	1
			1	4
			1	2
66. Rhynchopsidæ			1	1
			9	39 + 2 = 4
38. Stercorariidæ .		696–699	2	4
69. Procellariidæ			15	29 + 2 = 3
70. Podicipitidæ		729-735	5	7 + 1 =
71. Colymbidæ		736–740	1	5
72. Alcidæ		741-764	14	24 + 3 = 2

### SUMMARY.

Number of genera.	379
Number of species	764
Number of subspecies	160

### CONCORDANCE.

	CONCORDANCE,										
No. of old cat- alogue,	No. of new cat- alogue.	No. of old cat-	No. of new cat- alogue.	No. of old cat- alogue,	No. of new cat- alogue.	No. of old cat- alogue.	No. of new cat- alogue.	No. of old cat- alogue.	No. of new cat- alogue.	No. of old cat- alogue.	No. of new cat- alogue.
1	454	35	427	69	387	102	336	135	315	167	74
$\frac{1}{2}$	453	36	428	70	388	103	340	136	316	167a	74a
$\tilde{3}$	455	37	429	71	383	104	339	137	318	168	88
4	_	38	430	72	359	105	338	138	321	169	75
5	414	39	449	73	_	106	337	139	320	170	122
6.	414	40	_	74	360	107	349	140	325a	171	_
7	417	41	451	75	360b	108	350	141	325	172	120
8	415	42	452	76	361	109	351	142	326	173	121
9	419	43	451	77	361a	110	352	143	324	174	118
10	413	44	425	78	364	111	353	144	322	175	119
11	412	45	423	79	363	112	354	144a	323	176	123
12	412a	46	434	80	362	113	355	145	327	177	123a
13	420	47	394	81	366	114	357	146	328	178	77
14	433	48	405	82	367	115	357a	147	330	179	76
15	431	49	402	83	368	116	358	148	1	180	79
16	431	50	402b		-368a	116a	356	149	5b	181	81
17	432	51	395	85	369	117	382	149a	5a	182	78
18	442	. 52	396	86	369a	118	383	150	5	183	85
19	442	53	399	87	369b		381	151	2	183a	84
20	436b	54	397	88	370	120	333	152	4	184	86
21	442	55	400	89	370	121	332	153	4a	185	87
22	438	56	401	90	371	122	302	154	3	186	115
23	436	57	401	91	372	123	301	155	7	187	116
24	436b		408	92	373	124	304	156	9	188	117
25	439	59	408	93	374	125	303	157	21	189	107
26	439a		409	94	375	126	306	158	55	190	109
27	443	61	406	95	377	127	307	159	23	191	108
28	442	65	407	96	376	128	305	160	24	192	105
29	437	63	392	97	378	129	010	161	30	193	94
30	447	64	391	98	378b		312	162	33	194	95
31	447	65	384	98a	378a		313	163	32	195	96
32	448	66	389	99	379	132	311	164	19	196	102
33	445	67	389	100		133	314	165	71	197	100
34	426	68	385	101	335	134	317	166	73	198	111

### Concordance—Continued.

No. of old cat-	No. of	No. of old cat-	No. of	No. of	No. of	No. of	No. of	No. of	No. of	No. of	No. of
old cat- alogue.	new cat- alogue.	old cat- alogue.	new cat- alogue,	No. of old cat- alague,	new cat-	old cat- alogue.	new cat- alogue.	old cat- alogue.	new cat- alogue.	old cat-	new cat-
					aroguer	urogues.	arogue.	alogue.	atogue.	alogue.	alogue.
4400	443	350		0.4.3							
199	112	259	15	318	172	378	254	440	295	500	503
$\frac{200}{201}$	99 98	$259a \\ 260$	15 13a	$\frac{318a}{319}$	172a 173	379 380	255	441	294	500a	504
205	101	261	13a	320	179	381	244 245	442	296 297	501	505
203	93	261a	13	321	178	382	248	444	288	502 503	515
204	97	252	55	322	175	383	247	445	456	504	516
205	110	263	59a	323	174	384	251	446	457	505	523
206	90	234	58	324		385	250	447	458	503	522
207	91	265	60	325	186	386	249	448	459	507	517
208 209	113	256	60a	326	187	387	248	449	462	508	520
210	$\frac{103}{114}$	268	61 67	328	188 189	388 389	252 243	$\frac{450}{451}$	464	509	521
211	124	269	68	329	189	390	244	451	$\frac{460}{466}$	510 511	513 511
212	126	270	63	330	190	391	237	453	465	512	507
213	125	271	63a	331	191	392	238b	454	467	513	. 508
214	127	272	63	332	193a	393	238	455	468	514	_
215	127	273	65	333	193	394	238a	456-	469	515	509
216	132	274	35	334	194	395	241	457	470a	516	510
217 218	128 129	275 276	55 55a	335 336	193 <i>b</i> 196	393 397	$\frac{240b}{240}$	$\frac{458}{459}$	470	517	566
219	130	277	55a 51	337	197	398	239	460	471 472	518 519	567 565
220	161	278	51a	338	198	399	257	461	472a	520	564
221	164	279	52	339	199	400	258	462	479	521	563
222	163	280	53	340	200	401	261	463	478a	522	525
223	162	281 282	54	341	201	402	261a	464	477	523	526a
224 225	160 154	283	27 28	342 343	$202 \ 231e$	403	262	465	473	524	527
226	153	284	29	344	204	405	260	$\frac{465a}{466}$	$473a \ 473b$	525 526	527 <i>a</i> 529
227	155	285	36	345	206	406	263	437	474	527	535
258	156	283	37	346	207a	407	264	468	475	528	530
229	157	287	38	347	208	408	265	469	476	529	540
230	158	288	39	348	205	409	266	470	475	530	539a
$\frac{231}{231a}$	$\frac{152}{152a}$	289 289a	41a 41a	349 350	209 222	410 411	268	471 472	$\frac{480}{480b}$	531 532	534 538
232	150	290	41	351	221	412	267	473	481	533	536
233	151	291	41b	352	218	413	269	474	482	534	542
234	26	292	43	353	220	414	270	475	483	535	541
235	25	293	42	354	217	415	271	476	484	536	528
236	148	294 295	40	355	224	416	272	477	485	537	552
237 238	149 149a	296	$\frac{46}{45}$	356 357	$\frac{225}{210}$	417 418	$\frac{273}{274}$	478 479	582   583	538 539	$\frac{547}{548}$
239	-	297	49	358	214	419	275	480	584	540	549
240	135	298	47	359	211	420	277	481	581	541	550
241	136	299	48	360	212	421	278	482	491	542	553
242	100	300	50	361	213	422	278u	483	491	543	557
243 244	137 138	$\frac{301}{302}$	$\frac{159}{300}$	362 363	215 331	423 424	$\frac{280}{280}$	484 485	492 490	544 545	554 - 555
244	139	303	165	364	231b	425	281	485	489	546	อออ 556
246	145	304	166	365	231c	496	282	486a	489	547	553
247	142	305	168	366	231d	427	282a	487	487	548	545
248	143	306	168a	367	231a	428	282b	488	456	549	558
249	144	307	169	368	234	429	283	489	486	550	559
$\frac{250}{251}$	$\frac{141}{141a}$	308 209	170	369 · 370	233 226	430 431	$284 \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \$	490 491	493 498	551 552	560 569
252	140	301	184	371	228	432	286	492	497	553	571
253	11	311	_	372	230	433	287	493	494	554	572
253a	11	312		373	236	434	289	494	- 1	555	574
254	12	313	181	374	235	435	290	495	495	556	576
255 256	10	314 315	182	375 376	235a	436	290c	496 497	496   500	557 558	575 577
256 257	16 16a	316	182a 183	376a	235e $235b$	437 438	293 292	497	502	559	580
258	17	317	185	377	256	439	291	499	501	560	579

# 242 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.

## Concordance—Continued.

No. of old cat- alogue.	No. of new cat- alogue.	No. of old cat- alogue.	No. of new cat- alogue.	No. of old cat- alogue-	No. of new cat- alogue.	No. of old cat- alogue.	No. of new cat- alogue.	No. of old cat- alogue	No. of new cat- alogue.	No. of old cat- alogue.	No. of new cat- alogue.
561	578	589	615	620	642	651	706	681	679	710	741
561a	588	590	616	621	648	652	696	682	680	711	742
562	589	591	618	622	643b	653	697	683	681	712	745
563	591	592	617	623	643	654	698	684	682	713	744
563a	591a	593	620	624	643a	655	699	685	683	714	7436
564	590	594	619	625	644	656	660	686	685	715	743
565	593a	595	621	626	645	657	662	687	684	716	745
566	593a	596	622	627	646	658	661	688	681	717	746
567	594	597	623	628	649	659	662	689	686	718	746
567a	594c	598	625	629	654	660	663	690	687	719	748
568	594b	599	626	630	~01	661	664a	691	685	720	748
569	594a	600	624	631	701	662	664	692	688	721	749
570	595	601	632	632	702	663	668	693	687	722	750
571	596	602	633	633	703	664	669	694	690	723	750
572	597	603	633	634	704	664a	670	695	693	724	751
573	598	604	630	635	705	665	670 672	696	695	725	747
574	599	605	632	636	705a	666	673	697 698	656	726	760 761
575	600	606	627a	637	706	667	674		736	727	
576	601	607	628	638	717	668		699	738	728	762
577 578	602 605	608 609	629 634	639 640	719 726	669 670	674 675	700 701	· 739 740	729 730	763 763
579 580	612 611	610 611	635 636	641 642	727 723	671 672	658	702 703	731	731 732	764 755
581	609	612	637	643	723 724	673	674	703a	731	732	755 755
582	610	613	638	644	724	674	659	7034	731 729	734	759
583	608	614		645	721	675	659	704	739	735	756
584	604	615	640	646	721	676	657	706	730	736	753
585	607	616	641	647	709	677	657	707	733a	737	754
586	606	617	650	648	714	678	676	707	733	738	754 752
587	613	618	652	649	714	679	678	708a	734	100	132
588	614	619	639	650	712	680	677	709	735		

## INDEX TO THE GENERA.

[The numbers refer to the first species of each genus in the catalogue.]

	Page.		Page.
A.	4074	Basilinna	. 347
Accipiter	431	Berniela	. 594
Actodromas	533	Bonasa	. 473
Æchmophorus	729	Botaurus	. 497
Ægialitis	517	Brachyrhamphus	. 755
Ægiothus	178	Bubo	. 405
Æsalon	416	Budytes	
Agelæus	261	Buteo	
Aix	613	Butorides	
Ajaja	505	C.	
Alanda	299	Calamospiza	. 256
Alea	741	Calidris	
Alle	752	Callipepla	
Aluco	394	Calothorax	
Amazilia	345	Calypte	
Ammodromus	201	Campephilus	
Ampelis	150	Camptolemus	
Amphispiza	224	Campylorhynchus	
Anas	601	Canace	-
Anorthura	65	Caprimulgus	
Anous	695	Cardellina	
Anser	593		
Antenor	434	Cardinalis	
Anthus	71	Carpodacus	
Antrostomus	353	Catharista	
Aphelocoma	291	Cathartes	
Aphriza	511	Catherpes	
Aquila	449	Centrocercus	
Aramus	581	Centronyx	
Archibuteo	447	Centrophanes	
Ardea	486	Centurus	
Ardetta	498	Ceratorhina	
Arquatella	530	Certhia	
Asio	395	Certhiola	
Astragalinus	181	Ceryle	
Astur	433	Chætura	
Asturina	445	Chamea	
Atthis.	342	Chamæpelia	
Auriparus	50	Charadrius	201
Aythya	617	Chaulelasmus	
11, 111	011	Chen	004
В.		Chondestes	43.00.00
Bartramia	555	Chordeiles	
Basileuterus	133	Chrysomitris	184
		243	

	Page.		Page.
Ciceronia	750	G.	
Cinclus	19	Galeoscoptes	12
Circus	430	Gallinago	520
Cistothorus	68	Gallinula	579
Clangula	619	Garzetta	490
Coccyzus	386 .	Geococcyx	385
Colaptes	378	Geothlypis	120
Columba	456	Geotrygon	467
Colymbus	736	Glaucidium	409
Contopus	318	Grus	582
	392	Guiraea	246
Conurus		Gymnocitta	285
Corvus	250		
Cotile	157	H.	= 0
Coturniculus	198	Hæmatopus	500
Creagrus	678	Hadrostomus	333
Crex	577	Haliæëtus	451
Crotophaga	389	Halocyptena	720
Cupidonia	477	Harelda	623
Cyanecula	20	Harporhynchus	13
Cyanocitta	289	Helminthophaga	78
Cymochorea	723	Helmitherus	77
Cypseloides	350	Helon:ea	76
Cypselus	349	Herodias	489
Cyrtonyx	485	Hesperiphona	165
		Hesperocichla	9
Dafila	605	Heteroscelus	550
Daption	719	Hierofalco	418
Dendrocygna	599	Himantopus	567
· -	93	Hirundo	154
Dendræea		Histrionicus	629
Dichromanassa	491	Hydranassa	492
Diomedea	700	Hydrochelidon	693
Dolichonyx	257	Hylocichla	1
Dytes	732		371
E.		Hylotomus	911
Ectopistes	459	I.	
Engyptila	463	Iache	348
Elanoïdes	426	Ieteria	123
Elanus	427	Icterus	265
Embernagra	236	Ictinia	428
Empidonax	324	Ionornis	578
Eremophila	300		
Ereunetes	541	J.	
Ergatiens	132	Juneo	216
Erismatura	634	L,	
Eudocimus	501		474
Eugenes	334	Lagopus	626
Euphonia	160	Lampronetta	148
	100	Lanius	
F.	414	Lanivireo	140
Falco	414	Larus	660
Florida	493	Leucosticte	174
Fratercula	743	Limosa	543
Fregetta	728	Lobipes	564
Fulica	580	Lomvia	764
Fulix	614	Lophodytes	638
12-1	705	Lophophanes	36

	Page.		Dogo
Lophortyx	482	Р.	Page.
Loxia	172	Pachyrhamphus	332
Lunda	745	Pagophila	
	* 10	Pandion	
M.		Parra	
Machetes	554	Parula	
Macrorhamphus	527	Parus	
Mareca	606	Passerculus	
Megalestris	696	Passerella	
Melauerpes	375	Passerina	
Melanetta	631	Pediœcetes	
Meleagris	470		
Melopelia	464	Pelecanus	
Melospiza	231	Pelidna	
Mergus	636	Peliouetta	
Merula	7	Perisoreus	
Micrathene	411	Perissoglossa	
Micropalama	528	Petrochelidon	
	301	Peucæa	
Milyulus		Peucedramus	
Minus	11	Phaëthon	654
Mulotilta	74	Phainopepla	26
Molothrus	258	Phalacrocorax	642
Momotus	381	Phalænoptilus	355
Motacilla	69	Phalaropus	563
Myeteria	499	Phaleris	747
Myiadestes	25	Philaete	598
Myiarchus	311	Philohela	525
Myiodinastes	310	Phæbetria	703
Myiozetetes	309	Phænicopterus	
N.		Phonipara	
N.	~	Phylloseopus	34
Neocorys	73	Piea	286
Nettion.	611	Pieieorvus	284
Nomonyx	635	Picoides	
Numenius	558	Pieus	
Nyctale	400	Pinicola	166
Nyetea	407	Pipilo	
Nyetherodius	496	Pitangus	000
Nyetiardea	495	Pleetrophanes	4 30
Nyctidromus	356	_	503
0,		Plegadis	210
Oceanites	722		
Oceanodroma	726	Podasocys	731
Ochthodromus	522	Podilymbus	
Œdemia			2.00
	630	Polioptila	
Œstrelata	717	Polyborus	625
Olor	584	Polystieta	
Onychotes	446	Poweetes	
Operators	118	Porzana	706
Oreortyx	481	Priocella	707
Oreoscoptes	10	Priofinus	
Ornithion	331	Procellaria	721
Ortalis	469	Progne	152
Ortyx	480	Protonotaria	75
Ossifraga	704	Psaltriparus	47
Oxyechus	516	Pseudogryphus	-453

# 246 PROCEEDINGS OF UNITED STATES NATIONAL MUSEUM.

	Page.		Page.
Psilorhinus	288	Strix	397
Ptycorhamphus	751	Sturnella	263
Puffinus	708	Sturnus	279
Pyranga	161	Sula	650
Pyrrhula	167	Surnia	407
Pyrrhuloxia	243	Symphemia	559
Pyrocephalus	330	Synthliborhamphus	753
Q.		Т.	
Querquedula	609	Tachybaptes	734
Quiscalus	275	Tachycineta	155
R.		Tachypetes	639
Rallus	569	Tantalus	500
Recurvirostra	566	Telmatodytes	67
Regulus	30	Thrasaëtus	450
Rhodostethia	676	Thryomanes	61
		Thryothorus	60
Rhyacophilus	550	Tinnunculus	420
Rhynchofalco	419	Totanus	547
Rhynchophanes	190	Tringa	529
Rhynchops	656	Tringoides	557
Rhynchopsitta	391	Trochilus	335
Rissa	658		63
Rostrhamus	4.9	Troglodytes	384
s.		Trogon	
Salpinetes	58	Tryngites	556
Saxicola	21	Turdus	6
Sayornis	315	Tyrannus	303
Scardafella	466	U.	
Scolecophagus	273		
Scolopax	524	Ulula	399
Score		Uria	760
Scops	402	Urubitinga	444
Selasphorus	339	Utamania	742
Setophaga	128	V.	
Sialia	22	Vanellus	512
Simorhynchus	748		142
Sitta	51	Vireo	
Siurus	115	Vireosylvia	135
Somateria	627	W.	
Spatula	608		بالجالد
Spectyto	408	Wilsonia	124
Spermophila	252		
Sphyrapieus	369	X.	
Spiza	254	Xanthocephalus	260
Spizella	210	Xanthura	296
Squatarola	513	Xema	677
Starnænas	468	Xenopieus	366
Steganopus	565	g g	
Stelgidopteryx	158	Z.	
Stellula	343	Zamelodia	244
Stercorarius	697	Zenæda	462
Sterna	679	Zenædura	460
Strepsilas	509	Zonotrichia	205